arXiv:1508.04182v1 [math.RT] 18 Aug 2015

Categorifying the tensor product of the Kirillov-Reshetikhin crystal $B^{1,1}$ and a fundamental crystal

Henry Kvinge and Monica Vazirani

August 19, 2015

Abstract

We use Khovanov-Lauda-Rouquier (KLR) algebras to categorify a crystal isomorphism between a fundamental crystal and the tensor product of a Kirillov-Reshetikhin crystal and another fundamental crystal, all in affine type. The nodes of the Kirillov-Reshetikhin crystal correspond to a family of "trivial" modules. The nodes of the fundamental crystal correspond to simple modules of the corresponding cyclotomic KLR algebra. The crystal operators correspond to socle of restriction and behave compatibly with the rule for tensor product of crystal graphs.

Contents

1 Introduction **2**

2 Background and notation **3**
- 2.1 Cartan datum . . . 3
- 2.2 Review of crystals . . . 5
- 2.3 Perfect crystals and Kirillov-Reshetikhin crystals . . . 6
  - 2.3.1 Type A . . . 6
  - 2.3.2 General type . . . 9
  - 2.3.3 Paths on $B^{1,1}$ . . . 12

3 Key definitions: class $\mathcal{A}$, $\mathcal{B}$, $\mathcal{D}$ nodes and cyclotomic paths **14**

4 The KLR algebra R and some functors **16**
- 4.1 Definition of the KLR algebra $R(\nu)$ . . . 16
- 4.2 Induction, co-induction, and restriction . . . 18
- 4.3 Simple modules of $R(n\alpha_i)$ . . . 20
- 4.4 Crystal operators . . . 21
  - 4.4.1 The Serre relations . . . 24
- 4.5 Simple $R(c\alpha_i + \alpha_j)$-modules . . . 25
- 4.6 Jump . . . 25

5 Rep^Λ and the functor pr_Λ **29**
  - 5.0.1 Module-theoretic model of $B(\Lambda)$ . . . 31
  - 5.0.2 Interaction of pr_Λ and Ind . . . 31

6 The family of modules $T_{p;k}$ **34**

7 Main theorems **41**
7.1 Existence theorem 42
7.2 The action of the crystal operators 49

8 Appendix **54**
8.1 Simple modules in $\mathrm{Rep}^{\Lambda_h+\Lambda_i}$ in rank 2 54

1 Introduction

Kang-Kashiwara [11] and Webster [34] show the cyclotomic Khovanov-Lauda-Rouquier (KLR) algebra R^{Λ} categorifies the highest weight representation $V(\Lambda)$ in arbitrary symmetrizable type. (KLR algebras are also known in the literature as quiver Hecke algebras.) By a slight abuse of language, we will say the *combinatorial* version of this statement is that R^{Λ} categorifies the crystal $B(\Lambda)$, where simple modules correspond to nodes, and functors that take socle of restriction correspond to arrows, i.e. the Kashiwara crystal operators [23]. Webster [34] and Losev-Webster [25] categorify the tensor product of highest weight modules, and hence the tensor product of highest weight crystals. However, one can consider a tensor product of crystals

$$\mathcal{B} \otimes B(\Lambda) \simeq B(\Lambda') \tag{1}$$

where $\Lambda, \Lambda' \in P^+$ are of level k and $\mathcal{B}$ is a perfect crystal of level k. In this paper, we (combinatorially) categorify the crystal isomorphism (1) in the case $\Lambda = \Lambda_i$ is a fundamental weight and $\mathcal{B} = B^{1,1}$ is a Kirillov-Reshetikhin crystal. In other words, our main theorems give a purely module-theoretic construction of this crystal isomorphism. (One must modify the form of the crystal isomorphism in the case $B^{1,1}$ is not perfect or when Λ_i is not of level 1. See Section 2.3.2.) Each node of $B^{1,1}$ corresponds to an infinite family of "trivial" modules, but note this does *not* give a categorification of $\mathcal{B}$. These "trivial" modules $T_{p;k}$ are the KLR analogues of the nodes in highest weight crystals studied in [32] and are completely described in Section 6.

We note that this gives a construction of simple modules, starting from the $T_{p;k}$. In type A, this is somewhat intermediate between the crystal operator construction and the Specht module construction. See [33] for details. This paper also describes how socle of restriction interacts with the construction. One can also recover this for finite type whose Dynkin diagram is a subdiagram of that of type X_ℓ studied here. For a construction of simple modules related to the crystal $B(\infty)$ for finite type KLR algebras see [2].

This paper generalizes the theorems and constructions from [33] for type A affine, which were in turn originally proved for the affine Hecke algebra of type A at an $(\ell+1)$st root of unity [31, 28].

The authors welcome input on adding references if the ones included in this paper are incomplete.

Acknowledgments: We wish to thank Peter Tingley for interesting discussions and for pointing out we were using KR crystals and not just level 1 perfect crystals. The second author would like to thank David Hill for useful discussions and some initial jump computations.

The first author was partially supported by NSA grant H98230-12-1-0232, ICERM, and the Simons Foundation.

2 Background and notation

2.1 Cartan datum

We first review the Cartan datum associated with types $A_\ell^{(1)}$, $C_\ell^{(1)}$, $A_{2\ell}^{(2)}$, $A_{2\ell}^{(2)\dagger}$, $D_{\ell+1}^{(2)}$, $D_\ell^{(1)}$, $B_\ell^{(1)}$, and $A_{2\ell-1}^{(2)}$. Fix an integer $\ell \geq 2$. For each type X_ℓ listed above, $I = \{0, 1, \ldots, \ell\}$ will denote the indexing set. Let $[a_{ij}]_{i,j\in I}$ denote the associated Cartan matrix. We direct the reader to [9] for the explicit matrices. Following [9] we let $\mathfrak{h}$ be a Cartan subalgebra, $\prod = \{\alpha_0, \ldots, \alpha_\ell\}$ its system of simple roots, $\prod^\vee = \{h_0, \ldots, h_\ell\}$ its simple coroots, and Q and $Q^\vee$ the root and coroot lattices respectively. Then set

$$Q^+ = \bigoplus_{i\in I} \mathbb{Z}_{\geq 0}\alpha_i. \tag{2}$$

For an element $\nu \in Q^+$, we define its *height*, $|\nu|$, to be the sum of the coefficients, i.e. if $\nu = \sum_{i\in I} \nu_i \alpha_i$ then

$$|\nu| = \sum_{i\in I} \nu_i. \tag{3}$$

We also have a symmetric bilinear form

$$(\ ,\ ) : \mathfrak{h}^* \times \mathfrak{h}^* \to \mathbb{C}, \tag{4}$$

satisfying the property that

$$a_{ij} = \langle h_i, \alpha_j \rangle = \frac{2(\alpha_i, \alpha_j)}{(\alpha_i, \alpha_i)} \tag{5}$$

where $\langle\ ,\ \rangle : \mathfrak{h} \times \mathfrak{h}^* \to \mathbb{C}$ is the canonical pairing. Using this pairing we define the fundamental weights $\{\Lambda_i \mid i \in I\}$ via

$$\langle h_i, \Lambda_j \rangle = \delta_{ij}. \tag{6}$$

The weight lattice is $\bigoplus_{i\in I} \mathbb{Z}\Lambda_i$ and the integral dominant weights are

$$P^+ = \bigoplus_{i\in I} \mathbb{Z}_{\geq 0}\Lambda_i. \tag{7}$$

For all types considered in this paper, the associated Lie algebra $\mathfrak{g}$ has 1-dimensional center generated by the canonical central element $c = c_0 h_0 + \cdots + c_\ell h_\ell$ where $c_i \in \mathbb{Z}_{\geq 0}$. The level of a weight $\Lambda \in P^+$ is then defined to be $\langle c, \Lambda \rangle$.

For each type X_ℓ we draw below the associated Dynkin diagram and list the level of the fundamental weights.

- $A_1^{(1)}$, $I = \{0, 1\}$, Level 1 weights: Λ_0, Λ_1,



- $A_\ell^{(1)}$, $\ell \geq 2$, Level 1 weights: Λ_i, $i \in I$,



- $C_\ell^{(1)}$, $\ell \geq 2$, Level 1 weights: $\Lambda_i, i \in I$,



- $A_{2\ell}^{(2)}$, $\ell \geq 2$, Level 1 weight: Λ_0, Level 2 weights: $\Lambda_1, \ldots, \Lambda_\ell$



- $A_{2\ell}^{(2)\dagger}$, $\ell \geq 2$, Level 1 weight: Λ_ℓ, Level 2 weights: $\Lambda_0, \ldots, \Lambda_{\ell-1}$



- $D_{\ell+1}^{(2)}$, $\ell \geq 2$, Level 1 weights: Λ_0, Λ_ℓ, Level 2 weights: $\Lambda_1, \ldots, \Lambda_{\ell-1}$



- $D_\ell^{(1)}$, $\ell \geq 4$, Level 1 weights: Λ_0, Λ_1, $\Lambda_{\ell-1}$, Λ_ℓ, Level 2 weights: $\Lambda_2, \ldots, \Lambda_{\ell-2}$



- $B_\ell^{(1)}$, $\ell \geq 3$, Level 1 weights: Λ_0, Λ_1, Λ_ℓ, Level 2 weights: $\Lambda_2, \ldots, \Lambda_{\ell-1}$



- $A^{(2)}_{2\ell-1}$, $\ell \geq 3$, Level 1 weights: Λ_0, Λ_1, Level 2 weights: $\Lambda_2, \dots, \Lambda_\ell$



In most of the following theorems, we omit type $A^{(1)}_1$. All theorems and constructions in this paper hold in this type but require special arguments. These are presented in [33].

2.2 Review of crystals

We recall the tensor category of crystals following Kashiwara [16], see also [15, 14, 17].

A *crystal* is a set B together with maps

- $\mathrm{wt}\colon B \longrightarrow P$,
- $\varepsilon_i, \varphi_i : B \longrightarrow \mathbb{Z} \sqcup \{-\infty\}$ for $i \in I$,
- $\widetilde{e}_i, \widetilde{f}_i \colon B \longrightarrow B \sqcup \{0\}$ for $i \in I$,

such that

C1. $\varphi_i(b) = \varepsilon_i(b) + \langle h_i, \mathrm{wt}(b)\rangle$ for any i.

C2. If $b \in B$ satisfies $\widetilde{e}_i b \neq 0$, then

$$\varepsilon_i(\widetilde{e}_i b) = \varepsilon_i(b) - 1, \qquad \varphi_i(\widetilde{e}_i b) = \varphi_i(b) + 1, \qquad \mathrm{wt}(\widetilde{e}_i b) = \mathrm{wt}(b) + \alpha_i. \tag{8}$$

C3. If $b \in B$ satisfies $\widetilde{f}_i b \neq 0$, then

$$\varepsilon_i(\widetilde{f}_i b) = \varepsilon_i(b) + 1, \qquad \varphi_i(\widetilde{f}_i b) = \varphi_i(b) - 1, \qquad \mathrm{wt}(\widetilde{f}_i b) = \mathrm{wt}(b) - \alpha_i. \tag{9}$$

C4. For b_1, $b_2 \in B$, $b_2 = \widetilde{f}_i b_1$ if and only if $\widetilde{e}_i b_2 = b_1$.

C5. If $\varphi_i(b) = -\infty$, then $\widetilde{e}_i b = \widetilde{f}_i b = 0$.

For $b \in B$ we also define

$$\varepsilon(b) = \sum_{i \in I} \varepsilon_i(b)\Lambda_i \tag{10}$$

and

$$\varphi(b) = \sum_{i \in I} \varphi_i(b)\Lambda_i. \tag{11}$$

If B_1 and B_2 are two crystals, then a *morphism* $\psi\colon B_1 \to B_2$ of crystals is a map

$$\psi\colon B_1 \sqcup \{0\} \to B_2 \sqcup \{0\}$$

satisfying the following properties:

M1. $\psi(0) = 0$.

M2. If $\psi(b) \neq 0$ for $b \in B_1$, then

$$\mathrm{wt}(\psi(b)) = \mathrm{wt}(b), \qquad \varepsilon_i(\psi(b)) = \varepsilon_i(b), \qquad \varphi_i(\psi(b)) = \varphi_i(b). \tag{12}$$

M3. For $b \in B_1$ such that $\psi(b) \neq 0$ and $\psi(\widetilde{e}_i b) \neq 0$, we have $\psi(\widetilde{e}_i b) = \widetilde{e}_i(\psi(b))$.

M4. For $b \in B_1$ such that $\psi(b) \neq 0$ and $\psi(\widetilde{f}_i b) \neq 0$, we have $\psi(\widetilde{f}_i b) = \widetilde{f}_i(\psi(b))$.

A morphism ψ of crystals is called *strict* if

$$\psi \widetilde{e}_i = \widetilde{e}_i \psi, \qquad \psi \widetilde{f}_i = \widetilde{f}_i \psi, \tag{13}$$

and an *embedding* if ψ is injective.

Given two crystals B_1 and B_2 their tensor product $B_1 \otimes B_2$ (using the reverse Kashiwara convention) has underlying set $\{b_1 \otimes b_2 \mid b_1 \in B_1 \text{ and } b_2 \in B_2\}$ where we identify $b_1 \otimes 0 = 0 \otimes b_2 = 0$. The crystal structure is given as follows:

$$\mathrm{wt}(b_1 \otimes b_2) = \mathrm{wt}(b_1) + \mathrm{wt}(b_2), \tag{14}$$

$$\varepsilon_i(b_1 \otimes b_2) = \max\{\varepsilon_i(b_2), \varepsilon_i(b_1) - \langle h_i, \mathrm{wt}(b_2)\rangle\}, \tag{15}$$

$$\varphi_i(b_1 \otimes b_2) = \max\{\varphi_i(b_2) + \langle h_i, \mathrm{wt}(b_1)\rangle, \varphi_i(b_1)\}, \tag{16}$$

$$\widetilde{e}_i(b_1 \otimes b_2) = \begin{cases} \widetilde{e}_i b_1 \otimes b_2 & \text{if } \varepsilon_i(b_1) > \varphi_i(b_2) \\ b_1 \otimes \widetilde{e}_i b_2 & \text{if } \varepsilon_i(b_1) \leq \varphi_i(b_2), \end{cases} \tag{17}$$

$$\widetilde{f}_i(b_1 \otimes b_2) = \begin{cases} \widetilde{f}_i b_1 \otimes b_2 & \text{if } \varepsilon_i(b_1) \geq \varphi_i(b_2) \\ b_1 \otimes \widetilde{f}_i b_2 & \text{if } \varepsilon_i(b_1) < \varphi_i(b_2). \end{cases} \tag{18}$$

Given a crystal B, we can draw its associated crystal graph with nodes (or vertices) B and I-colored arrows (directed edges) as follows. When $\widetilde{e}_i b = a$ (so $b = \widetilde{f}_i a$) we draw an i-colored arrow $a \xrightarrow{i} b$. We also say b has an incoming i-arrow and a has an outgoing i-arrow.

2.3 Perfect crystals and Kirillov-Reshetikhin crystals

2.3.1 Type A

In type $A_\ell^{(1)}$, the highest weight crystal $B(\Lambda_i)$ has a model (see Figure 1) with nodes the $(\ell+1)$-*restricted* partitions, i.e. $\lambda = (\lambda_1, \ldots, \lambda_t)$ such that $\lambda_r \in \mathbb{Z}_{\geq 0}$, $0 \leq \lambda_r - \lambda_{r+1} < \ell + 1$ for all r. Let $B^{1,1}$ be the crystal graph



$B^{1,1}$ is also drawn in (29) without the node labels.

$B^{1,1}$ is an example of a *perfect* crystal (see [13] for the definition and important properties). One key property this level 1 perfect crystal has is that tensoring it with a fundamental (or highest weight level 1) crystal yields an isomorphism to another level 1 highest weight crystal. In particular, for $i \in I$ there exists an isomorphism of crystals,

$$\mathcal{T} : B(\Lambda_i) \xrightarrow{\simeq} B^{1,1} \otimes B(\Lambda_{i-1}).$$

The isomorphism is pictured in Figure 2 for $i = 0$ and $\ell = 2$. Note the underlying graph of $B(\Lambda_i)$ is identical to that of $B(\Lambda_0)$, but the colors of the arrows are obtained from those of $B(\Lambda_0)$ by adding i mod $(\ell+1)$.



Figure 1: $B(\Lambda_0)$ and $B(\Lambda_2)$ for $\ell = 2$.

Combinatorially, $\mathcal{T}(\lambda) = \boxed{k} \otimes \mu$ where $k \equiv \lambda_1 + i - 1 \bmod (\ell + 1)$ and $\mu = (\lambda_2, \ldots, \lambda_t)$ if $\lambda = (\lambda_1, \lambda_2, \cdots, \lambda_t)$. So we obtain μ from λ by removing its top row. In Figure 2, we draw

$$\mathcal{T}(\lambda) = \begin{array}{c} \boxed{k} \\ \otimes \\ \mu \end{array} \tag{19}$$

so the visual of the top row removal stands out. Note $\lambda_1 - \mu_1 < \ell + 1$ means that $\mathcal{T}$ has a well-defined inverse.

When drawing $B(\Lambda_i)$, we label each box of a partition with $k \in I$, such that the main diagonal gets label i, and labels increase by 1 mod $(\ell + 1)$ as one increases diagonals (moving right). In this manner, the last box in the top row of λ is labeled k when $\mathcal{T}(\lambda) = \textcircled{k} \otimes \mu$. Note further that if we have a k-arrow $\gamma \xrightarrow{k} \lambda$ then the box λ/γ is labeled k (though not necessarily conversely). This box is often called a "good" k-box. In fact, once one knows the structure of $B^{1,1}$ and the tensor product rule for crystals, one can obtain the rule for which k-box $\widetilde{e}_k$ removes by iterating $\mathcal{T}$.



Figure 2: The isomorphism $B(\Lambda_0) \simeq B^{1,1} \otimes B(\Lambda_2)$ for $\ell = 2$.

Remark 2.1. There is another model of $B(\Lambda_i)$ for which nodes are indexed by $(\ell+1)$-regular partitions. A partition λ is $(\ell+1)$-*regular* iff its transpose λ^T is $(\ell+1)$-restricted. (However one does not obtain this model by merely transposing the partition indexing each node.) There is a similar isomorphism $\mathcal{T}' : B(\Lambda_i) \xrightarrow{\simeq} B^{\ell,1} \otimes B(\Lambda_{i+1})$ corresponding to column removal.

For type $A_\ell^{(1)}$ the diagram for $B^{\ell,1}$ is



(20)

This can be obtained from $B^{1,1}$ in type $A_\ell^{(1)}$ by reversing the direction of all arrows.

See Section 5.0.1 for another model of $B(\Lambda_i)$.

2.3.2 General type

The perfect crystal $B^{1,1}$ in (29) is also an example of a Kirillov-Reshetikhin (KR) crystal. For a quantized affine algebra $U_q'(\mathfrak{g})$, the KR crystals $B^{r,s}$ correspond to a special family of finite dimensional modules $W^{r,s}$ indexed by a positive integer s and a Dynkin node r from the classical subalgebra $\mathfrak{g}_0$ of $\mathfrak{g}$ [20], [27]. In this paper we work with the crystals $B^{1,1}$ which have particularly simple graphs. In all of the types we consider, with the exception of $C_\ell^{(1)}$, the crystal $B^{1,1}$ is perfect of level 1 [7]. When $B^{1,1}$ is perfect and Λ_i is a level 1 fundamental weight for $i \in I$, $B^{1,1}$ has a unique node b_i such that $\varepsilon(b_i) = \Lambda_i$ and $\varphi(b_i) = \Lambda_{\sigma(i)}$ for some $\sigma(i) \in I$. There then exists a crystal isomorphism [13]

$$\mathcal{T} : B(\Lambda_{\sigma(i)}) \xrightarrow{\sim} B^{1,1} \otimes B(\Lambda_i). \tag{21}$$

In this paper we will also consider the tensor product $B^{1,1} \otimes B(\Lambda_i)$ when Λ_i is a level 2 weight and consider type $C_\ell^{(1)}$ where $B^{1,1}$ is not perfect. In those cases, $B^{1,1} \otimes B(\Lambda_i)$ decomposes as

$$B^{1,1} \otimes B(\Lambda_i) \cong \bigoplus_{j \in I} k_j B(s_j \Lambda_j), \qquad k_j \in \mathbb{Z}_{\geq 0}, s_j \in \{1, 2\} \tag{22}$$

for appropriate k_j, s_j. See [12], [10] for the case where Λ_i is not level 1, and [24] for type $C_\ell^{(1)}$, $i = 0$ which can be generalized to $i \in I$ by similar methods. (When the level of Λ_i is 1, all $s_j = 1$; but when the level of Λ_i is 2 and the level of Λ_j is only 1, we can have $s_j = 2$.)

Example 2.2. In type $C_\ell^{(1)}$, one can check

$$B^{1,1} \otimes B(\Lambda_1) \simeq B(\Lambda_0) \oplus B(\Lambda_2). \tag{23}$$

Recall $B^{1,1}$ is *not* a perfect crystal in type $C_\ell^{(1)}$.

If we call the crystal isomorphism $\mathcal{T}$, label the nodes of $B^{1,1}$ as



(24)

and the highest weight node of $B(\Lambda_i)$ as b_{Λ_i} then

$$\mathcal{T}(\boxed{1} \otimes b_{\Lambda_1}) = b_{\Lambda_2} \tag{25}$$

$$\mathcal{T}(\boxed{\bar{1}} \otimes b_{\Lambda_1}) = b_{\Lambda_0} \tag{26}$$

as $\varepsilon(\boxed{1}) = \varepsilon(\boxed{\bar{1}}) = \Lambda_1$ but $\varphi(\boxed{1}) = \Lambda_2, \varphi(\boxed{\bar{1}}) = \Lambda_0$.

How this difference manifests in the main theorems of the paper is also addressed in Remark 7.9 and Example 7.16.

Note, one can also check $B^{1,1} \otimes B(\Lambda_0) \simeq B(\Lambda_1)$.

In type $A_{2\ell}^{(2)}$ one can check that for similar reasons as above that

$$B^{1,1} \otimes B(\Lambda_2) \simeq B(\Lambda_1) \oplus B(\Lambda_3), \tag{27}$$

if $\ell \geq 3$. Recall that $\Lambda_1, \Lambda_2, \Lambda_3$ are all of level 2, whereas Λ_0 is level 1. One can also check

$$B^{1,1} \otimes B(\Lambda_1) \simeq B(2\Lambda_0) \oplus B(\Lambda_2). \tag{28}$$

Below we draw the crystal graphs for the KR crystals $B^{1,1}$ for each type X_ℓ we are considering. The nodes in $B^{1,1}$ are labelled with ⬭ to distinguish them from the Dynkin nodes ○ . The conditions $\ell \geq 2, 3, 4$ hold below in each respective type as they do for the Dynkin diagrams listed in Section 2.1.

- $A_\ell^{(1)}$



(29)

- $C_\ell^{(1)}$



(30)

- $A_{2\ell}^{(2)}$



(31)

- $A_{2\ell}^{(2)\dagger}$



(32)

- $D_{\ell+1}^{(2)}$



(33)

- $D_{\ell}^{(1)}$



(34)

- $B_{\ell}^{(1)}$



(35)

- $A_{2\ell-1}^{(2)}$



(36)

2.3.3 Paths on $B^{1,1}$

Definition 2.3. Let X_ℓ be one of the affine types listed in Section 2.1 and let I be the indexing set of its Dynkin nodes. A type X_ℓ *path* p of length k, is a function $p : \{0, 1, \ldots, k-1\} \to I$ such that there is a directed walk in the type X_ℓ crystal $B^{1,1}$ whose ith step corresponds to a $p(i)$-arrow.

We specify the edge color data but not node data of the walk. There might then appear to be some danger of ambiguity of two different walks corresponding to the same path, as $B^{1,1}$ may have more than one arrow of a given color. Indeed, the single step paths $p : \{0\} \to I$ in general do not specify a unique walk in $B^{1,1}$. However, for paths with length k with $k \geq 2$, the definition above is sufficient to specify a unique walk.

Proposition 2.4. Let $p : \{0, 1, \ldots, k-1\} \to I$ be a type X_ℓ path of length k. If $k > 1$, then there is a unique walk in $B^{1,1}$ that corresponds to p.

Proof. This proposition is based on three observations about the graph $B^{1,1}$:

- A node a never has two incoming (respectively outgoing) arrows of the same color (this of course is true of all crystals, even when $\varphi_i(a) > 1$ or $\varepsilon_i(a) > 1$).
- There are at most two i-arrows in $B^{1,1}$ and hence at most two nodes a and b with an incoming (respectively outgoing) i-arrow.
- If a and b are nodes that each have an incoming (respectively outgoing) i-arrow, a has an outgoing (respectively incoming) j_1-arrow and b has an outgoing (respectively incoming) j_2-arrow, then $j_1 \neq j_2$.

By the first observation, a walk corresponding to p in $B^{1,1}$ is uniquely determined once we have chosen its initial node.

By the second observation, if $p(0) = i$ there are at most two choices (a and b) for the second node of the walk. If $k > 1$ then there is a second $p(1)$-arrow leaving either a or b. By the third observation, a and b cannot both have an outgoing $p(1)$-arrow. Assume without loss of generality that a has an outgoing $p(1)$-arrow. Then p uniquely corresponds to the walk in $B^{1,1}$ that starts by entering node a via a $p(0)$-arrow and leaves a by a $p(1)$-arrow. ☐

Because of the uniqueness implied by Proposition 2.4 we will henceforth use the terms "type X_ℓ path" and "walk in $B^{1,1}$" interchangeably when $k > 1$.

Example 2.5. Let $p : \{0,1,2\} \to I$ be the path of length 3 in type $D_5^{(1)}$, defined such that $p(0) = 0$, $p(1) = 2$, $p(2) = 3$. We mark the walk corresponding to p with a dotted line.



(37)

Definition 2.6. For a path $p : \{0,1,\dots,k-1\} \to I$ we call the arrow corresponding to $p(0)$ the *tail* of p, and the arrow corresponding to $p(k-1)$ the *head* of p. An extension to the tail of p by a j-arrow is a path $p' : \{0,1,\dots,k\} \to I$ such that $p'(t) = p(t-1)$ for $1 \le t \le k$ and $p'(0) = j$. An extension to the head of p by a j-arrow, is a path $p'' : \{0,1,\dots,k\} \to I$ such that $p''(t) = p(t)$ for $0 \le t \le k-1$ and $p''(k) = j$.

Let $\pi(j)$ be the length 1 path $\pi(j) : \{0\} \to I$, $\pi(j)(0) = j$. For a path p, we denote the extension of its tail by a j-arrow by $\pi(j) \star p$ and the extension of its head by a j-arrow by $p \star \pi(j)$. We can think of extension as concatenation of paths. If the tail (respectively head) of p cannot be extended by a j-arrow then we set $\pi(j) \star p = \mathbf{0}$ (respectively $p \star \pi(j) = \mathbf{0}$).

The set of colors of arrows that can extend the tail of a path p of length $k > 1$ is denoted ext_p^-,

$$\mathrm{ext}_p^- := \{\ j \mid \pi(j) \star p \neq \mathbf{0}\ \}. \tag{38}$$

The set of colors of arrows that can extend the head of p is denoted ext_p^+,

$$\mathrm{ext}_p^+ := \{\ j \mid p \star \pi(j) \neq \mathbf{0}\ \}. \tag{39}$$

When ext_p^- (respectively ext_p^+) contains a single element i, we allow ourselves the convenience of writing $p(-1) = i$ (respectively $p(k) = i$).

Example 2.7. In example 2.5

$$\mathrm{ext}_p^+ = \{4,5\} \qquad \mathrm{ext}_p^- = \{1\}. \tag{40}$$

so we write $p(-1) = 1$.

Remark 2.8. In general, when p has length 1, then ext_p^- and ext_p^+ are not well-defined. However, the symmetry of the $B^{1,1}$ graphs means that $\mathrm{ext}_p^- \cup \mathrm{ext}_p^+$ is well-defined even for length 1 paths.

3 Key definitions: class $\mathcal{A}$, $\mathcal{B}$, $\mathcal{D}$ nodes and cyclotomic paths

Let $i, j, j', a, a', b, b' \in I$. We classify the arrows of $B^{1,1}$ into three classes: $\mathcal{A}$, $\mathcal{B}$, and $\mathcal{D}$.

- An i-arrow belongs to class $\mathcal{B}$ if it is adjacent to another i-arrow. That is, it either enters a node that has an outgoing i-arrow or it departs from a node with an incoming i-arrow. For instance, in the diagram below both i-arrows are of class $\mathcal{B}$.



- A pair of arrows a and a' are said to be a class $\mathcal{D}$ pair if they are the first and second step respectively of a bifurcation (a bifurcation occurs at a node of degree 3). Below both (a, a') and (b, b') are class $\mathcal{D}$ pairs.



For all $B^{1,1}$ it will actually be the case that $a = b'$, $a' = b$, and $j = j'$, whenever (a, a') is a class $\mathcal{D}$ pair.



- All other arrows in $B^{1,1}$ are class $\mathcal{A}$. For large ℓ, the vast majority of arrows will belong to class $\mathcal{A}$.

Remark 3.1. It is not hard to see from inspection that within a given type, our classification of arrows in $B^{1,1}$ descends to a classification of elements of I in that type. In other words within a $B^{1,1}$ diagram, there are never two i-arrows that belong to different classes. So it makes sense to talk about $i \in I$ being of a particular class. Table 1 classifies the elements of I for all types X_ℓ considered in this paper.

Below is one of our key definitions.

Definition 3.2. Let p be a path of length $k \geq 1$ in $B^{1,1}$. p is called a *cyclotomic path of tail weight* $(\Lambda_{i_1}, \Lambda_{i_2})$ if the following hold:

1. $p(0) = i_2$ and $p(1) \neq i_2$. If p has length 1 then $p \star \pi(i_2) = \mathbf{0}$, i.e. the head of p cannot be extended by an i_2-arrow.

Type	class $\mathcal{A}$	class $\mathcal{B}$	class $\mathcal{D}$ pairs
$A_\ell^{(1)}$	$0, 1, \ldots, \ell-1, \ell$		
$C_\ell^{(1)}$	$0, 1, \ldots, \ell-1, \ell$		
$A_{2\ell}^{(2)}$	$1, \ldots, \ell-1, \ell$	0	
$A_{2\ell}^{(2)\dagger}$	$0, 1, \ldots, \ell-1$	ℓ	
$D_{\ell+1}^{(2)}$	$1, \ldots, \ell-1$	$0, \ell$	
$D_\ell^{(1)}$	$2, 3, \ldots, \ell-2$		$(0, 1), (\ell-1, \ell)$
$B_\ell^{(1)}$	$2, \ldots, \ell-1$	ℓ	$(0, 1)$
$A_{2\ell-1}^{(2)}$	$2, \ldots, \ell-1, \ell$		$(0, 1)$

Table 1: Classification of elements of I for each type X_ℓ

2. $p(0)$ and $p(1)$ are not a class $\mathcal{D}$ pair. If p has length 1 then there is no $i \in I$ such that the head of p can be extended by an i-arrow and $(p(0), i)$ is a class $\mathcal{D}$ pair.

3. $\text{ext}_p^- = \{i_1\}$, i.e. p has a unique extension by an i_1-arrow to its tail.

4. $(\pi(i_1) \star (\pi(i_1) \star p)) = \mathbf{0}$, i.e. p cannot have its tail extended twice by i_1-arrows.

The following proposition follows from inspection of the $B^{1,1}$ crystal graphs.

Proposition 3.3. 1. Let X_ℓ be any type not equal to $D_3^{(2)}$, $D_4^{(1)}$ and $B_3^{(1)}$. Then for all $i \in I$ there exists some $j \in I$ such that there is a cyclotomic path p of tail weight (Λ_j, Λ_i).

2. In type $D_3^{(2)}$ so long as $i \neq 1$, i.e. $i \in \{0, 2\}$, there is $j \in I$ such that there exists a cyclotomic path p of tail weight (Λ_j, Λ_i).

3. In type $D_4^{(1)}$ and $B_3^{(1)}$ so long as $i \neq 2$, i.e. $i \in \{0, 1, 3\}$, there is some $j \in I$ such that there exists a cyclotomic path p of tail weight (Λ_j, Λ_i).

When the above paths exists they can be of any length $k \in \mathbb{Z}_{\geq 1}$.

Definition 3.4. We refer to $1 \in I$ in type $D_3^{(2)}$ and $2 \in I$ in types $D_4^{(1)}$ and $B_3^{(1)}$ as *forbidden* elements of I.

Remark 3.5. Above we saw that the one-to-one correspondence between length k paths in $B^{1,1}$ and length k directed walks in $B^{1,1}$ breaks down when $k = 1$. It is important to note that this does not occur for length 1 cyclotomic paths because of the extra information imposed by the unique tail extension.

4 The KLR algebra R and some functors

4.1 Definition of the KLR algebra $R(\nu)$

Fix an indeterminant q. In what follows we let $q_i = q^{\frac{(\alpha_i,\alpha_i)}{2}}$,

$$[k]_i = q_i^{k-1} + q_i^{k-3} + \cdots + q_i^{1-k} \quad \text{and} \quad [k]_i! = [k]_i[k-1]_i \dots [1]_i. \tag{41}$$

For $\nu = \sum_{i\in I} \nu_i\alpha_i$ in Q^+ with $|\nu| = m$, we define $\mathrm{Seq}(\nu)$ to be all sequences

$$\underline{\mathbf{i}} = (i_1, i_2, \dots, i_m) \tag{42}$$

such that i_k appears ν_k times. For $\underline{\mathbf{i}} \in \mathrm{Seq}(\nu)$ and $\underline{\mathbf{j}} \in \mathrm{Seq}(\mu)$, $\underline{\mathbf{ij}}$, will denote the concatenation of the two sequences unless otherwise specified. It follows that $\underline{\mathbf{ij}} \in \mathrm{Seq}(\nu + \mu)$. We write

$$i^n = (\underbrace{i, i, \dots, i}_{n}). \tag{43}$$

There is a left action of the symmetric group, $\mathcal{S}_m$, on $\mathrm{Seq}(\nu)$ defined by,

$$s_k(\underline{\mathbf{i}}) = s_k\Big(i_1, i_2, \dots, i_k, i_{k+1}, \dots, i_m\Big) = (i_1, i_2, \dots, i_{k+1}, i_k, \dots, i_m) \tag{44}$$

where s_k is the adjacent transposition in $\mathcal{S}_m$ that interchanges k and $k+1$.

Our presentation of KLR algebras below follows that found in [18] and [19]. Using the more general definition with Rouquier's parameters $Q_{i,j}(u,v)$ will not change the results or proofs in this paper, as they concern crystal-theoretic phenomena. There is also a useful diagrammatic presentation of KLR algebras which can be found in [18], [19]. By results of Brundan-Kleshchev [4], [5], there is an isomorphism between the cyclotomic KLR algebra, $R^\Lambda(\nu)$, of type $A_\ell^{(1)}$, and H_ν^Λ where H_ν^Λ is a block of the cyclotomic Hecke algebra H_m^Λ as defined in [1, 3, 6]. Hence readers unfamiliar with KLR algebras can translate all statements and proofs for the type $A_\ell^{(1)}$, $\Lambda = \Lambda_i$ case in terms of Hecke algebras throughout the paper (historically, this is the original setting in which the theorems from this paper were proved). In fact for type $A_\ell^{(1)}$ the reader can think of all results as being stated for $\mathbb{F}_{\ell+1}\mathcal{S}_m$ in the case that $\ell + 1$ is prime.

For $\nu \in Q^+$ with $|\nu| = m$, the *KLR algebra $R(\nu)$* associated with Cartan matrix $[a_{ij}]_{i,j\in I}$ is the associative, graded, unital $\mathbb{C}$-algebra generated by

$$1_{\underline{\mathbf{i}}} \ \text{for} \ \underline{\mathbf{i}} \in \mathrm{Seq}(\nu), \quad x_r \ \text{for} \ 1 \le r \le m, \quad \psi_r \ \text{for} \ 1 \le r \le m-1, \tag{45}$$

subject to the following relations, where $\underline{\mathbf{i}}, \underline{\mathbf{j}} \in \mathrm{Seq}(\nu)$:

$$1_{\underline{\mathbf{i}}}1_{\underline{\mathbf{j}}} = \delta_{\underline{\mathbf{i}},\underline{\mathbf{j}}}1_{\underline{\mathbf{i}}}, \qquad x_r 1_{\underline{\mathbf{i}}} = 1_{\underline{\mathbf{i}}} x_r, \qquad \psi_r 1_{\underline{\mathbf{i}}} = 1_{s_r(\underline{\mathbf{i}})}\psi_r, \qquad x_r x_t = x_t x_r, \tag{46}$$

$$\psi_r\psi_t = \psi_t\psi_r \qquad \text{if } |r-t|>1, \tag{47}$$

$$\psi_r\psi_r 1_{\underline{\mathbf{i}}} = \begin{cases} 0 & \text{if } i_r = i_{r+1}, \\ 1_{\underline{\mathbf{i}}} & \text{if } (\alpha_{i_r}, \alpha_{i_{r+1}}) = 0, \\ (x_r^{-a_{i_r i_{r+1}}} + x_{r+1}^{-a_{i_{r+1} i_r}})1_{\underline{\mathbf{i}}} & \text{if } (\alpha_{i_r}, \alpha_{i_{r+1}}) \neq 0 \text{ and } i_r \neq i_{r+1}, \end{cases} \tag{48}$$

$$(\psi_r\psi_{r+1}\psi_r - \psi_{r+1}\psi_r\psi_{r+1})1_{\underline{\mathbf{i}}} = \begin{cases} \displaystyle\sum_{t=0}^{-a_{i_r i_{r+1}}-1} x_r^t x_{r+2}^{-a_{i_r i_{r+1}}-1-t} 1_{\underline{\mathbf{i}}} & \text{if } i_r = i_{r+2} \text{ and } (\alpha_{i_r}, \alpha_{i_{r+1}}) \neq 0, \\ 0 & \text{otherwise,} \end{cases} \tag{49}$$

$$(\psi_r x_t - x_{s_r(t)}\psi_r)1_{\underline{\mathbf{i}}} = \begin{cases} 1_{\underline{\mathbf{i}}} & \text{if } t = r \text{ and } i_r = i_{r+1}, \\ -1_{\underline{\mathbf{i}}} & \text{if } t = r+1 \text{ and } i_r = i_{r+1}, \\ 0 & \text{otherwise.} \end{cases} \tag{50}$$

The elements $1_{\underline{\mathbf{i}}}$ are idempotents in $R(\nu)$ by (46) and the identity element is

$$1_\nu = \sum_{\underline{\mathbf{i}} \in \mathrm{Seq}(\nu)} 1_{\underline{\mathbf{i}}}. \tag{51}$$

Thus, as a vector space $R(\nu)$ decomposes as,

$$R(\nu) = \bigoplus_{\underline{\mathbf{i}},\underline{\mathbf{j}} \in \mathrm{Seq}(\nu)} 1_{\underline{\mathbf{i}}} R(\nu) 1_{\underline{\mathbf{j}}}. \tag{52}$$

The generators of $R(\nu)$ are graded via

$$\deg(1_{\underline{\mathbf{i}}}) = 0, \quad \deg(x_r 1_{\underline{\mathbf{i}}}) = (\alpha_{i_r}, \alpha_{i_r}), \quad \deg(\psi_r 1_{\underline{\mathbf{i}}}) = -(\alpha_{i_r}, \alpha_{i_{r+1}}). \tag{53}$$

We define

$$R = \bigoplus_{\nu \in Q^+} R(\nu). \tag{54}$$

Notice that while $R(\nu)$ is unital, R is not.

For each $w \in \mathcal{S}_m$ we fix once and for all a reduced expression

$$\widehat{w} = s_{i_1} s_{i_2} \dots s_{i_t}. \tag{55}$$

Here the s_k are Coxeter generators of $\mathcal{S}_m$, and t is the Coxeter length. Let $\psi_{\widehat{w}} = \psi_{i_1}\psi_{i_2}\dots\psi_{i_t}$ correspond to the chosen reduced expression $\widehat{w}$ of w. For $\underline{\mathbf{i}}, \underline{\mathbf{j}} \in \mathrm{Seq}(\nu)$, let ${}_{\underline{\mathbf{j}}}\mathcal{S}_{\underline{\mathbf{i}}}$ be the permutations in $\mathcal{S}_m$ that take $\underline{\mathbf{i}}$ to $\underline{\mathbf{j}}$.

Theorem 4.1. [18, Theorem 2.5] As a $\mathbb{C}$-vector space $1_{\underline{\mathbf{j}}} R(\nu) 1_{\underline{\mathbf{i}}}$ has basis,

$$\{\psi_{\widehat{w}} x_1^{b_1} \dots x_m^{b_m} 1_{\underline{\mathbf{i}}} \mid w \in {}_{\underline{\mathbf{j}}}\mathcal{S}_{\underline{\mathbf{i}}},\ b_r \in \mathbb{Z}_{\geq 0}\}. \tag{56}$$

It is known that all simple $R(\nu)$-modules are finite dimensional [18]. For this reason, in this paper we only consider the category of finite-dimensional KLR-modules denoted by $R(\nu)\text{-}\mathrm{mod}$ and $R\text{-}\mathrm{mod}$. We write $G_0(R)$ for the Grothendieck group.

We often refer to $1_{\underline{\mathbf{i}}}M$ as the $\underline{\mathbf{i}}$-weight space of M and any $0 \neq v \in 1_{\underline{\mathbf{i}}}M$ as a weight vector. A weight basis is a basis consisting of weight vectors.

We define the graded character of an $R(\nu)$-module to be

$$\mathrm{Char}(M) = \sum_{\underline{\mathbf{i}} \in \mathrm{Seq}(\nu)} \mathrm{gdim}(1_{\underline{\mathbf{i}}}M) \cdot [\underline{\mathbf{i}}]. \tag{57}$$

Here $\mathrm{gdim}(1_{\underline{\mathbf{i}}}M)$ is an element of $\mathbb{Z}[q, q^{-1}]$, and hence $\mathrm{Char}(M)$ is an element of the free $\mathbb{Z}[q, q^{-1}]$-module generated by all $[\underline{\mathbf{i}}]$ for $\underline{\mathbf{i}} \in \mathrm{Seq}(\nu)$. We will let $\mathrm{supp}(M)$ denote the multiset that is the support of $\mathrm{Char}(M)|_{q=1}$ so that

$$\mathrm{Char}(M)|_{q=1} = \sum_{[\underline{\mathbf{i}}] \in \mathrm{supp}(M)} [\underline{\mathbf{i}}]. \tag{58}$$

Our notational convention is to write $\underline{\mathbf{i}} \in \mathrm{Seq}(\nu)$ but write $[\underline{\mathbf{i}}] \in \mathrm{supp}(M)$. Since characters are an important combinatorial tool, it is worthwhile to set a special notation for them.

Remark 4.2. A simple module is determined by its character, hence $[M] \in G_0(R)$ the Grothendieck group, is also determined by its character [18], [19].

Because R is a graded algebra, we will only work with homomorphisms between R-modules that are either degree preserving or degree homogeneous. We denote the $\mathbb{C}$-vector space of degree preserving homomorphisms between $R(\nu)$-modules M and N by $\mathrm{Hom}(M, N)$. Since any homogeneous homomorphism can be interpreted as degree preserving by shifting the grading on our target or source module, we can write the $\mathbb{C}$-vector space of homogeneous homomorphisms between M and N, $\mathrm{HOM}(M, N)$, by

$$\mathrm{HOM}(M, N) = \bigoplus_{k \in \mathbb{Z}} \mathrm{Hom}(M, N\{k\}). \tag{59}$$

While the grading is important, it is shown in [18] that there is a unique grading on a simple R-module up to overall grading shift. Since this paper concerns simple modules, we will rarely use or discuss the grading. All isomorphisms between modules will be taken up to overall grading shift.

Remark 4.3. Because $x_r 1_{\underline{\mathbf{i}}} \in R(\nu)$ is always positively graded for $1 \leq r \leq |\nu|$ and $\underline{\mathbf{i}} \in \mathrm{Seq}(\nu)$, then on a finite dimensional $R(\nu)$-module, M, $x_r 1_{\underline{\mathbf{i}}}$ will always act nilpotently.

4.2 Induction, co-induction, and restriction

It was shown in [18] and [19] that for $\nu, \mu \in Q^+$ there is a non-unital embedding

$$R(\nu) \otimes R(\mu) \hookrightarrow R(\nu + \mu). \tag{60}$$

This map sends the idempotent $1_{\underline{\mathbf{i}}} \otimes 1_{\underline{\mathbf{j}}}$ to $1_{\underline{\mathbf{ij}}}$. The identity $1_\nu \otimes 1_\mu$ of $R(\nu) \otimes R(\mu)$ has as its image

$$\sum_{\underline{\mathbf{i}} \in \mathrm{Seq}(\nu)} \sum_{\underline{\mathbf{j}} \in \mathrm{Seq}(\mu)} 1_{\underline{\mathbf{ij}}}. \tag{61}$$

Using this embedding one can define induction and restriction functors,

$$\begin{aligned} \mathrm{Ind}_{\nu,\mu}^{\nu+\mu} : (R(\nu)\otimes R(\mu))\text{-}\mathrm{mod} &\to R(\nu+\mu)\text{-}\mathrm{mod} && (62)\\ M &\mapsto R(\nu+\mu)\otimes_{R(\nu)\otimes R(\mu)} M && (63)\end{aligned}$$

and

$$\mathrm{Res}_{\nu,\mu}^{\nu+\mu} : R(\nu+\mu)\text{-}\mathrm{mod} \to (R(\nu)\otimes R(\mu))\text{-}\mathrm{mod}. \qquad (64)$$

In the future we will write $\mathrm{Ind}_{\nu,\mu}^{\nu+\mu} = \mathrm{Ind}$ and $\mathrm{Res}_{\nu,\mu}^{\nu+\mu} = \mathrm{Res}$ when the algebras are understood from the context. More generally we can extend this embedding to finite tensor products

$$R(\nu^{(1)})\otimes R(\nu^{(2)})\otimes\cdots\otimes R(\nu^{(k)}) \hookrightarrow R(\nu^{(1)}+\nu^{(2)}+\cdots+\nu^{(k)}). \qquad (65)$$

We refer to the image of this embedding as a *parabolic subalgebra* and denote it by $R(\underline{\nu}) \subset R(\nu^{(1)}+\cdots+\nu^{(k)})$. We denote the image of the identity under this embedding as $1_{\underline{\nu}}$. It follows from Theorem 4.1 that $R(\nu^{(1)}+\nu^{(2)}+\cdots+\nu^{(k)})1_{\underline{\nu}}$ is a free right $R(\underline{\nu})$-module, and $1_{\underline{\nu}}R(\nu^{(1)}+\nu^{(2)}+\cdots+\nu^{(k)})$ is a free left $R(\underline{\nu})$-module.

Let $m_i = |\nu^{(i)}|$ and set

$$P = (m_1,\dots,m_k) \quad\text{and}\quad \mathcal{S}_P = \mathcal{S}_{m_1}\times\mathcal{S}_{m_1}\times\cdots\times\mathcal{S}_{m_k}. \qquad (66)$$

Let $\mathcal{S}_{m_1+\cdots+m_k}/\mathcal{S}_P$ be the collection of minimal length left coset representatives of $\mathcal{S}_P$ in $\mathcal{S}_{m_1+\cdots+m_k}$ and $\mathcal{S}_P\backslash\mathcal{S}_{m_1+\cdots+m_k}$ be the collection of minimal length right coset representatives of $\mathcal{S}_P$ in $\mathcal{S}_{m_1+\cdots+m_k}$. We can construct a weight basis for an induced module as follows. If M is an $R(\underline{\nu})$-module with weight basis U then $\mathrm{Ind}_{\underline{\nu}}^{\nu^{(1)}+\cdots+\nu^{(k)}} M$ has weight basis

$$\{\psi_{\widehat{w}}\otimes u \mid w\in\mathcal{S}_{m_1+\cdots+m_k}/\mathcal{S}_P, u\in U\}. \qquad (67)$$

Induction is left adjoint to restriction (a property known as Frobenius reciprocity),

$$\mathrm{HOM}_{R(\nu^{(1)}+\cdots+\nu^{(k)})}(\mathrm{Ind}_{\underline{\nu}}^{\nu^{(1)}+\cdots+\nu^{(k)}} M, N) \cong \mathrm{HOM}_{R(\underline{\nu})}(M, \mathrm{Res}_{\underline{\nu}}^{\nu^{(1)}+\cdots+\nu^{(k)}} N). \qquad (68)$$

The right adjoint to restriction is the co-induction functor coInd: $R(\underline{\nu})\text{-}\mathrm{mod} \to R(\nu^{(1)}+\cdots+\nu^{(k)})\text{-}\mathrm{mod}$,

$$\mathrm{coInd}_{\underline{\nu}}^{\nu^{(1)}+\cdots+\nu^{(k)}} - := \mathrm{HOM}_{R(\underline{\nu})}(R(\nu^{(1)}+\cdots+\nu^{(k)}), -), \qquad (69)$$

so there is an isomorphism of $\mathbb{C}$-vectors spaces

$$\mathrm{HOM}_{R(\nu^{(1)}+\cdots+\nu^{(k)})}(N, \mathrm{coInd}_{\underline{\nu}}^{\nu^{(1)}+\cdots+\nu^{(k)}} M) \cong \mathrm{HOM}_{R(\underline{\nu})}(\mathrm{Res}_{\underline{\nu}}^{\nu^{(1)}+\cdots+\nu^{(k)}} N, M). \qquad (70)$$

For KLR algebras there exists a useful connection between induced and co-induced modules.

Proposition 4.4. [23] Let M be a finite dimensional $R(\mu)$-module and N a finite dimensional $R(\nu)$-module. Then up to grading shift

$$\mathrm{Ind}_{\mu,\nu}^{\mu+\nu} M\boxtimes N \cong \mathrm{coInd}_{\nu,\mu}^{\mu+\nu} N\boxtimes M. \qquad (71)$$

This proposition immediately tells us that most properties that we can prove about induced modules can be transferred to co-induced modules with appropriate modifications.

Remark 4.5. Suppose we have

$$\operatorname{Ind} M \boxtimes D \overset{f}{\twoheadrightarrow} A \tag{72}$$

with A a simple $R(\mu{+}\nu)$-module, D an $R(\nu)$-module, and M an $R(\mu)$-module with a filtration

$$0 = M_0 \subseteq M_1 \subseteq \cdots \subseteq M_t = M. \tag{73}$$

Consider the restriction

$$\operatorname{Ind} M_i \boxtimes D \xrightarrow{f} A \tag{74}$$

and suppose this is nonzero, hence surjective. Suppose further that $f(\operatorname{Ind} M_{i-1} \boxtimes D) = 0$; then we get a nonzero map

$$\operatorname{Ind} M_i/M_{i-1} \boxtimes D \xrightarrow{f} A. \tag{75}$$

By abuse of notation we will say *f factors through* $\operatorname{Ind} M_i/M_{i-1} \boxtimes D$ in this case (note we have extended the usual notation of factoring through M/M_{i-1} to M_i/M_{i-1}).

Given $\underline{\mathbf{i}} \in \operatorname{Seq}(\nu)$ and $\underline{\mathbf{j}} \in \operatorname{Seq}(\mu)$, a shuffle of $\underline{\mathbf{i}}$ and $\underline{\mathbf{j}}$ is an element $\underline{\mathbf{k}}$ of $\operatorname{Seq}(\nu + \mu)$ such that $\underline{\mathbf{k}}$ has $\underline{\mathbf{i}}$ as a subsequence and $\underline{\mathbf{j}}$ as the complementary subsequence. We denote by $\underline{\mathbf{i}} \sqcup \underline{\mathbf{j}}$ the formal sum of all shuffles of $\underline{\mathbf{i}}$ and $\underline{\mathbf{j}}$. The multi-set of all shuffles of $\underline{\mathbf{i}}$ and $\underline{\mathbf{j}}$ are in bijection with the minimal length left coset representatives of $\mathcal{S}_{|\nu|+|\mu|}/\mathcal{S}_{|\nu|} \times \mathcal{S}_{|\mu|}$. Using the definition of degree from KLR algebras, we can associate to any shuffle a degree which we denote as $\deg(\underline{\mathbf{i}}, \underline{\mathbf{j}}, \underline{\mathbf{k}})$. Then the quantum shuffle of $\underline{\mathbf{i}}$ and $\underline{\mathbf{j}}$ is

$$\underline{\mathbf{i}} \mathbin{\text{ш}} \underline{\mathbf{j}} = \sum_{\sigma \in \mathcal{S}_{|\nu|+|\mu|}/\mathcal{S}_{|\nu|} \times \mathcal{S}_{|\mu|}} q^{\deg(\underline{\mathbf{i}}, \underline{\mathbf{j}}, \sigma(\underline{\mathbf{ij}}))} \sigma(\underline{\mathbf{ij}}), \tag{76}$$

so that $\underline{\mathbf{i}} \sqcup \underline{\mathbf{j}} = (\underline{\mathbf{i}} \mathbin{\text{ш}} \underline{\mathbf{j}})|_{q=1}$. Note that we will usually shuffle characters, hence we write $[\underline{\mathbf{i}}] \sqcup [\underline{\mathbf{j}}]$. For an $R(\mu)$-module M and $R(\nu)$-module N it was shown in [18] that

$$\operatorname{Char}(\operatorname{Ind}_{\mu,\nu}^{\mu+\nu} M \boxtimes N) = \operatorname{Char}(M) \mathbin{\text{ш}} \operatorname{Char}(N). \tag{77}$$

This identity is referred to as the **Shuffle Lemma**.

4.3 Simple modules of $R(n\alpha_i)$

For $\nu = n\alpha_i$, induction allows for a particularly easy description of all simple $R(n\alpha_i)$-modules. Let $L(i)$ be the 1-dimensional $R(\alpha_i)$-module where $x_1 1_i$ acts as zero. Then up to overall grading shift the unique simple $R(n\alpha_i)$-module is

$$L(i^n) = \operatorname{Ind}_{\alpha_i, \alpha_i, \ldots, \alpha_i}^{n\alpha_i} L(i) \boxtimes \cdots \boxtimes L(i). \tag{78}$$

Some authors prefer to shift the degree so that the character is

$$\operatorname{Char}(L(i^n)) = [n]_i! [i, i, \ldots, i]. \tag{79}$$

4.4 Crystal operators

In the previous section we defined induction and restriction for KLR algebras. Following the work of Grojnowski [8] where crystal operators were developed as functors on the category of modules over affine Hecke algebras of type A (or [22] for $\mathbb{F}_{\ell+1}\mathcal{S}_m$), the KLR analogues of crystal operators were introduced in [18], and further developed in [23], [11]. For each $i \in I$, if $M \in R(\nu)$ - mod and $\nu - \alpha_i \in Q^+$, define the functor $\Delta_i : R(\nu)\text{ - mod} \to R(\nu-\alpha_i) \otimes R(\alpha_i)\text{ - mod}$ as the restriction

$$\Delta_i M := \operatorname{Res}^{\nu}_{\nu-\alpha_i,\alpha_i} M \tag{80}$$

Note that this is equivalent to multiplying M by $1_{\nu-\alpha_i} \otimes 1_{\alpha_i}$. It is also sometimes useful to think of this functor as killing all weight spaces corresponding to elements of $\operatorname{Seq}(\nu)$ that do not end in i. If $\nu - \alpha_i \notin Q^+$ then $\Delta_i M = \mathbf{0}$. We similarly define

$$\Delta_{i^n} M := \operatorname{Res}^{\nu}_{\nu-n\alpha_i,n\alpha_i} M. \tag{81}$$

Next define the functor $e_i : R(\nu)\text{ - mod} \to R(\nu - \alpha_i)\text{ - mod}$ as the restriction,

$$e_i M := \operatorname{Res}^{R(\nu-\alpha_i)\otimes R(\alpha_i)}_{R(\nu-\alpha_i)} \circ \Delta_i M, \tag{82}$$

hence it is an exact functor. When M is simple we can further refine this functor by setting

$$\widetilde{e}_i M := \operatorname{soc} e_i M. \tag{83}$$

We measure how many times we can apply $\widetilde{e}_i$ to M by

$$\varepsilon_i(M) := \max\{n \geq 0 \mid (\widetilde{e}_i)^n M \neq \mathbf{0}\ \}. \tag{84}$$

Let $\widetilde{f}_i : R(\nu)\text{ - mod} \to R(\nu + \alpha_i)\text{ - mod}$ be defined by

$$\widetilde{f}_i M := \operatorname{cosoc} \operatorname{Ind} M \boxtimes L(i) \tag{85}$$

(still assuming M is simple). Some of the most important facts about $e_i, \widetilde{e}, \widetilde{f}_i$ stated in [18] are given in the following proposition

Proposition 4.6. Let $i \in I$, $\nu \in Q^+$, $n \in \mathbb{Z}_{>0}$.

1. Let $M \in R(\nu)$ - mod. Then

$$\operatorname{Char}(\Delta_{i^n} M) = \sum_{\underline{\mathbf{j}} \in \operatorname{Seq}(\nu-n\alpha_i)} \operatorname{gdim}(1_{\underline{\mathbf{j}} i^n} M) \cdot \underline{\mathbf{j}} i^n,$$

2. Let $N \in R(\nu)$ - mod be irreducible and $M = \operatorname{Ind}^{\nu+n\alpha_i}_{\nu,n\alpha_i} N \boxtimes L(i^n)$. Let $\varepsilon = \varepsilon_i(N)$. Then
   - (a) $\Delta_{i^{\varepsilon+n}} M \cong (\widetilde{e}_i)^{\varepsilon} N \boxtimes L(i^{\varepsilon+n})$.
   - (b) $\operatorname{cosoc} M$ is irreducible, and $\operatorname{cosoc} M \cong (\widetilde{f}_i)^n N$, $\Delta_{i^{\varepsilon+n}} (\widetilde{f}_i)^n N \cong (\widetilde{e}_i)^{\varepsilon} N \boxtimes L(i^{\varepsilon+n})$, and $\varepsilon_i((\widetilde{f}_i)^n N) = \varepsilon + n$.
   - (c) $(\widetilde{f}_i)^n N$ occurs with multiplicity one as a composition factor of M.
   - (d) All other composition factors K of M have $\varepsilon_i(K) < \varepsilon + n$.

3. Let $\underline{\mu} = (\mu_1\alpha_i, \dots, \mu_r\alpha_i)$ with $\sum_{k=1}^{r} \mu_k = n$.
   (a) All composition factors of $\operatorname{Res}_{\underline{\mu}}^{n\alpha_i} L(i^n)$ are isomorphic to $L(i^{\mu_1}) \boxtimes \cdots \boxtimes L(i^{\mu_r})$, and $\operatorname{soc}(\operatorname{Res}_{\underline{\mu}}^{n\alpha_i} L(i^n))$ is irreducible.
   (b) $\widetilde{e}_i L(i^n) \cong L(i^{n-1})$.
4. Let $M \in R(\nu)$ - mod be irreducible with $\varepsilon_i(M) > 0$. Then $\widetilde{e}_i M = \operatorname{soc}(e_i M)$ is irreducible and $\varepsilon_i(\widetilde{e}_i M) = \varepsilon_i(M) - 1$. Furthermore if K is a composition factor of $e_i M$ and $K \ncong \widetilde{e}_i M$, then $\varepsilon_i(K) < \varepsilon_i(M) - 1$.
5. For irreducible $M \in R(\nu)$ - mod let $m = \varepsilon_i(M)$. Then $e_i^m M$ is isomorphic to $(\widetilde{e}_i)^m M^{\oplus [m]_i!}$. In particular, if $m = 1$ then $e_i M = \widetilde{e}_i M$.
6. For irreducible modules $N \in R(\nu)$ - mod and $M \in R(\nu + \alpha_i)$ - mod we have $\widetilde{f}_i N \cong M$ if and only if $N \cong \widetilde{e}_i M$.
7. Let $M, N \in R(\nu)$ - mod be irreducible. Then $\widetilde{f}_i M \cong \widetilde{f}_i N$ if and only if $M \cong N$. Assuming $\varepsilon_i(M), \varepsilon_i(N) > 0$, $\widetilde{e}_i M \cong \widetilde{e}_i N$ if and only if $M \cong N$.

On the level of characters, e_i roughly removes an i from the rightmost entry of a module's character. We can construct analogous functors for removal of i from the left side of a module's character, as well as an analogue to $\widetilde{f}_i$. These are denoted by $e_i^{\vee}$, $\widetilde{e}_i^{\vee}$, $\widetilde{f}_i^{\vee}$ and we will use them extensively in this paper. We use the involution σ introduced below to define them. Let w_0 be the longest element of $\mathcal{S}_{|\nu|}$. Then $\sigma : R(\nu) \to R(\nu)$ is defined as follows:

$$\begin{aligned} 1_{\underline{\mathbf{i}}} &\mapsto 1_{w_0(\underline{\mathbf{i}})} \\ x_r &\mapsto x_{|\nu|+1-r} \\ \psi_r 1_{\underline{\mathbf{i}}} &\mapsto (-1)^{\delta_{i_r, i_{r+1}}} \psi_{|\nu|-r} 1_{w_0(\underline{\mathbf{i}})}. \end{aligned}$$

For an $R(\nu)$-module M, let $\sigma^* M$ be the $R(\nu)$-module M but with the action of $R(\nu)$ twisted by σ,

$$r \cdot u = \sigma(r) u. \tag{86}$$

Now let $e_i^{\vee} : R(\nu)$ - mod $\to R(\nu - \alpha_i)$ - mod be the restriction functor defined as

$$e_i^{\vee} := \sigma^* e_i \sigma = \operatorname{Res}_{R(\nu-\alpha_i)}^{R(\alpha_i)\otimes R(\nu-\alpha_i)} \circ \operatorname{Res}_{\alpha_i, \nu-\alpha_i}^{\nu}, \tag{87}$$

and similarly,

$$\widetilde{e}_i^{\vee} M := \sigma^*(\widetilde{e}_i(\sigma^* M)) = \operatorname{soc} e_i^{\vee} M, \tag{88}$$

$$\widetilde{f}_i^{\vee} M := \sigma^*(\widetilde{f}_i(\sigma^* M)) = \operatorname{cosoc} \operatorname{Ind}_{\alpha_i, \nu}^{\nu+\alpha_i} L(i) \boxtimes M, \tag{89}$$

$$\varepsilon_i^{\vee}(M) := \varepsilon_i(\sigma^* M) = \max\{m \geq 0 \mid (\widetilde{e}_i^{\vee})^m M \neq \mathbf{0}\}. \tag{90}$$

It is important to note that by the exactness of restriction, $e_i, e_i^{\vee}$ are exact functors, while $\widetilde{e}_i$ and $\widetilde{e}_i^{\vee}$ are only left exact, and $\widetilde{f}_i$ and $\widetilde{f}_i^{\vee}$ are only right exact.

If $M \in R(\nu)$ - mod, we define $\operatorname{wt}(M) = -\nu$ and $\operatorname{wt}_i(M) = -\langle h_i, \nu \rangle$. The functors $\widetilde{e}_i$, $\widetilde{f}_i$, as well as ε_i and wt will be part of a crystal structure with simple R-modules as nodes [23].

Remark 4.7. There is a convenient character theoretic interpretation of ε_i and $\varepsilon_i^\vee$. Let M be a simple $R(\nu)$-module with $|\nu| = m$. Then

a.) $\varepsilon_i(M) = c$ implies that there exists

$$\underline{\mathbf{i}} = (i_1, \ldots, i_{m-c}, \underbrace{i, i, \ldots, i}_{c}) \tag{91}$$

such that $1_{\underline{\mathbf{i}}}A \neq \mathbf{0}$. In other words $[\underline{\mathbf{i}}]$ is in the support of M; however no $[\underline{\mathbf{j}}]$ such that

$$\underline{\mathbf{j}} = (i_1, \ldots, i_{m-c-1}, \underbrace{i, i, \ldots, i}_{c+1}). \tag{92}$$

is in $\mathrm{supp}(M)$.

b.) $\varepsilon_i^\vee(M) = c$ implies that there exists $[\underline{\mathbf{i}}]$ in the support of M of the form

$$\underline{\mathbf{i}} = (\underbrace{i, i, \ldots, i}_{c}, i_{c+1}, \ldots, i_m) \tag{93}$$

but no $[\underline{\mathbf{j}}]$ of the form

$$\underline{\mathbf{j}} = (\underbrace{i, i, \ldots, i}_{c+1}, i_{c+2}, \ldots, i_m). \tag{94}$$

Proposition 4.8. [30], [21] Suppose that M is a simple $R(\nu)$-module with $\varepsilon_i(M) > 0$. Then up to grading shift $\mathrm{cosoc}(e_i M) \cong \widetilde{e}_i M$.

The following proposition describes how the restriction functors e_i interact with induction.

Proposition 4.9. [29, Theorem 5.3] Let M be a simple $R(\mu)$-module and N be a simple $R(\nu)$-module. Then for $i \in I$ there is a short exact sequences

$$\mathbf{0} \to \mathrm{Ind}\, M \boxtimes e_i N \to e_i\Big(\mathrm{Ind}\, M \boxtimes N\Big) \to \mathrm{Ind}\, e_i M \boxtimes N \to \mathbf{0} \tag{95}$$

and

$$\mathbf{0} \to \mathrm{Ind}\, e_i^\vee M \boxtimes N \to e_i^\vee\Big(\mathrm{Ind}\, M \boxtimes N\Big) \to \mathrm{Ind}\, M \boxtimes e_i^\vee N \to \mathbf{0}. \tag{96}$$

Note that it follows from this result that if $e_i M = \mathbf{0}$ then

$$e_i\Big(\mathrm{Ind}\, M \boxtimes N\Big) \cong \mathrm{Ind}\, M \boxtimes e_i N \tag{97}$$

and similarly if $e_i N = \mathbf{0}$ then

$$e_i\Big(\mathrm{Ind}\, M \boxtimes N\Big) \cong \mathrm{Ind}\, e_i M \boxtimes N. \tag{98}$$

An analogous remark applies to $e_i^\vee$.

Proposition 4.10. Suppose that M is a simple $R(\mu)$-module, N is a simple $R(\nu)$-module, A is simple $R(\mu+\nu)$-module and there is a surjection

$$\mathrm{Ind}\, M \boxtimes N \twoheadrightarrow A. \tag{99}$$

1. If $\varepsilon_i(A) \neq 0$ then:
   (a) If $\varepsilon_i(M) = 0$ then there is a surjection
   $$\operatorname{Ind} M \boxtimes \widetilde{e}_i N \twoheadrightarrow \widetilde{e}_i A. \tag{100}$$
   (b) If $\varepsilon_i(N) = 0$, then there is a surjection
   $$\operatorname{Ind} \widetilde{e}_i M \boxtimes N \twoheadrightarrow \widetilde{e}_i A. \tag{101}$$
2. If $\varepsilon_i^\vee(A) \neq 0$ then:
   (a) If $\varepsilon_i^\vee(M) = 0$ there is a surjection
   $$\operatorname{Ind} M \boxtimes \widetilde{e}_i^\vee N \twoheadrightarrow \widetilde{e}_i^\vee A. \tag{102}$$
   (b) If $\varepsilon_i^\vee(N) = 0$ there is a surjection
   $$\operatorname{Ind} \widetilde{e}_i^\vee M \boxtimes N \twoheadrightarrow \widetilde{e}_i^\vee A. \tag{103}$$

Proof. We prove Part 1a, the other parts follow from analogous arguments. By Frobenius reciprocity it follows that $\varepsilon_i(N) \leq \varepsilon_i(A)$. By the exactness of the functor e_i and Proposition 4.9, $\varepsilon_i(N) \geq \varepsilon_i(A)$ so that $\varepsilon_i(N) = \varepsilon_i(A)$. Also by Proposition 4.9, the map of (99) induces a surjection
$$\operatorname{Ind} M \boxtimes e_i N \twoheadrightarrow e_i A. \tag{104}$$
Composing this with the surjection from Proposition 4.8 gives
$$\operatorname{Ind} M \boxtimes e_i N \twoheadrightarrow \widetilde{e}_i A. \tag{105}$$
From Remark 4.5 there exists a composition factor K of $e_i N$ such that there is a surjection,
$$\operatorname{Ind} M \boxtimes K \twoheadrightarrow \widetilde{e}_i A. \tag{106}$$
We must have
$$\varepsilon_i(K) = \varepsilon_i(\widetilde{e}_i A) = \varepsilon_i(A) - 1 = \varepsilon_i(N) - 1. \tag{107}$$
Proposition 4.6.4, then forces $K \cong \widetilde{e}_i N$. ☐

4.4.1 The Serre relations

Because the functors e_i, $i \in I$, are exact, they descend to well-defined linear operators on the Grothendieck group of R, $G_0(R)$. By abuse of notation, we will also call these operators e_i. We define divided powers
$$\begin{aligned} e_i^{(r)} : G_0(R) &\to G_0(R) \\ [M] &\mapsto \frac{1}{[r]_i!}[e_i^r M]. \end{aligned}$$

It is shown in [18, 19] that these operators satisfy the quantum Serre relations, and that these relations are in fact minimal. We have

$$\sum_{r=0}^{-a_{ij}+1} (-1)^r e_i^{(-a_{ij}+1-r)} e_j e_i^{(r)} [M] = \mathbf{0} \tag{108}$$

for all $i \neq j \in I$, and $M \in R\text{-mod}$. (Recall $a_{ij} = \langle h_i, \alpha_j \rangle$.) The minimality of these relations imply that, for $0 < c < -a_{ij} + 1$,

$$\sum_{r=0}^{c} (-1)^r e_i^{(c-r)} e_j e_i^{(r)} \tag{109}$$

is never the zero operator on $G_0(R)$ by the quantum Gabber-Kac Theorem [26] and the work of [18, 19], which essentially computes the kernel of the map from the free algebra on generators e_i to $G_0(R)$.

4.5 Simple $R(c\alpha_i + \alpha_j)$-modules

Assume that $i \neq j$ and set $a = a_{ij} = \langle h_i, \alpha_j \rangle$. We introduce the notation

$$\mathcal{L}(i^{c-n} j i^n) \tag{110}$$

for the irreducible $R(c\alpha_i + \alpha_j)$-modules when $c \leq -a$.

Theorem 4.11. [23] Let $c \leq -a$ and let $\nu = c\alpha_i + \alpha_j$. For each n with $0 \leq n \leq c$, there exists a unique irreducible $R(\nu)$-module denoted $\mathcal{L}(i^{c-n} j i^n)$ with

$$\varepsilon_i(\mathcal{L}(i^{c-n} j i^n)) = n. \tag{111}$$

Furthermore

$$\varepsilon_i^\vee(\mathcal{L}(i^{c-n} j i^n)) = c - n \tag{112}$$

and

$$\mathrm{Char}(\mathcal{L}(i^{c-n} j i^n)) = [c-n]_i! [n]_i! i^{c-n} j i^n. \tag{113}$$

Observe the support of $\mathcal{L}(i^{c-n} j i^n)$ is exactly $i^{c-n} j i^n$.

Proposition 4.12. [23] For $m > 0$,

$$\mathrm{Ind}\, \mathcal{L}(i^{-a-n} j i^n) \boxtimes L(i^m) \cong \mathrm{Ind}\, L(i^m) \boxtimes \mathcal{L}(i^{-a-n} j i^n) \tag{114}$$

is irreducible.

4.6 Jump

When we apply $\widetilde{f}_i$ to irreducible $R(\nu)$-module M for $i \in I$, then Proposition 4.6.2 tells us that $\widetilde{f}_i M$ is an irreducible $R(\nu + \alpha_i)$-module with

$$\varepsilon_i(\widetilde{f}_i M) = \varepsilon_i(M) + 1. \tag{115}$$

We could also ask whether $\varepsilon_i^\vee(\widetilde{f}_i M)$ and $\varepsilon_i^\vee(M)$ differ. Questions like this motivate the introduction of the function jump_i, which is based on a concept for Hecke algebras in [8], and was introduced for KLR algebras and studied extensively in [23].

Definition 4.13. Let M be a simple $R(\nu)$-module, and let $i \in I$. Then

$$\mathrm{jump}_i(M) := \max\{J \geq 0 \mid \varepsilon_i^\vee(M) = \varepsilon_i^\vee(\widetilde{f}_i^J M)\}. \tag{116}$$

Lemma 4.14. [23] Let M be a simple $R(\nu)$-module. The following are equivalent:

1. $\mathrm{jump}_i(M) = 0$
2. $\widetilde{f}_i M \cong \widetilde{f}_i^\vee M$
3. $\mathrm{Ind}\, M \boxtimes L(i^m)$ is irreducible for all $m \geq 1$
4. $\mathrm{Ind}\, M \boxtimes L(i^m) \cong \mathrm{Ind}\, L(i^m) \boxtimes M$ for all $m \geq 1$
5. $\mathrm{wt}_i(M) + \varepsilon_i(M) + \varepsilon_i^\vee(M) = 0$ (recall that $\mathrm{wt}_i(M) = -\langle h_i, \nu\rangle$),
6. $\varepsilon_i(\widetilde{f}_i^\vee M) = \varepsilon_i(M) + 1$
7. $\varepsilon_i^\vee(\widetilde{f}_i M) = \varepsilon_i^\vee(M) + 1$

Proof. See [23]. ☐

Example 4.15. For $c \leq -a_{ij}$ we can calculate $\mathrm{jump}_i(\mathcal{L}(i^{c-n} j i^n)) = -a_{ij} - c$. Proposition 4.12 follows from the equivalence of 1, 3, and 4 in Lemma 4.14.

It is shown in [23] that

$$\mathrm{jump}_i(\widetilde{f}_i M) = \max\{0, \mathrm{jump}_i(M) - 1\} = \mathrm{jump}_i(\widetilde{f}_i^\vee M). \tag{117}$$

This means that if we know $\mathrm{jump}_i(M)$, we can easily calculate $\mathrm{jump}_i(\widetilde{f}_i^k M)$ for any $k \geq 0$. It is also shown in [23] that

$$\mathrm{jump}_i(M) = \mathrm{wt}_i(M) + \varepsilon_i(M) + \varepsilon_i^\vee(M). \tag{118}$$

Using information from jump_i we can also determine when the crystal operators commute with their σ-symmetric versions.

Example 4.16. In type $A_\ell^{(1)}$ for $\ell > 1$, observe that $\mathrm{jump}_1(L(0)) = 1$ and

$$\widetilde{f}_1^\vee \widetilde{f}_1 L(0) \cong \mathrm{Ind}\, L(1) \boxtimes \mathcal{L}(01) \tag{119}$$

whose character has support $\{[1,0,1],[0,1,1],[0,1,1]\}$. However

$$\widetilde{f}_1 \widetilde{f}_1^\vee L(0) \cong \mathrm{Ind}\, \mathcal{L}(10) \boxtimes L(1) \tag{120}$$

whose character has support $\{[1,0,1],[1,1,0],[1,1,0]\}$.

In $A_1^{(1)}$, note that $\mathrm{jump}_1(\mathcal{L}(01)) = 1$ and we can similarly calculate $\widetilde{f}_1^\vee \widetilde{f}_1 \mathcal{L}(01) \not\cong \widetilde{f}_1 \widetilde{f}_1^\vee \mathcal{L}(01)$ (in fact the former is 8-dimensional while the latter is 4-dimensional).

We shall see below that this phenomenon is special to $\mathrm{jump}_i(M) = 1$.

Lemma 4.17. Let M be a simple $R(\nu)$-module.

1. [23] If $i \neq j$, then
   (a) $\widetilde{f}_i \widetilde{f}_j^\vee M \cong \widetilde{f}_j^\vee \widetilde{f}_i M$.
   (b) If $\widetilde{e}_j^\vee M \neq \mathbf{0}$ then $\widetilde{f}_i \widetilde{e}_j^\vee M \cong \widetilde{e}_j^\vee \widetilde{f}_i M$.
   (c) If $\widetilde{e}_j M \neq \mathbf{0}$ then $\widetilde{f}_i^\vee \widetilde{e}_j M \cong \widetilde{e}_j \widetilde{f}_i^\vee M$.
   (d) If further $\widetilde{e}_i M \neq \mathbf{0}$ then, $\widetilde{e}_i \widetilde{e}_j^\vee(M) \cong \widetilde{e}_j^\vee \widetilde{e}_i(M)$.
2. (a) $\mathrm{jump}_i(M) \neq 1$ if and only if $\widetilde{f}_i^\vee \widetilde{f}_i M \cong \widetilde{f}_i \widetilde{f}_i^\vee M$.
   (b) If $\widetilde{e}_i^\vee M \neq \mathbf{0}$, then $\mathrm{jump}_i(\widetilde{e}_i^\vee M) \neq 1$ if and only if $\widetilde{e}_i^\vee \widetilde{f}_i M \cong \widetilde{f}_i \widetilde{e}_i^\vee M$.
   (c) If $\widetilde{e}_i M \neq \mathbf{0}$, then $\mathrm{jump}_i(\widetilde{e}_i M) \neq 1$ if and only if $\widetilde{e}_i \widetilde{f}_i^\vee M \cong \widetilde{f}_i^\vee \widetilde{e}_i M$.

Proof.

1. Consider the short exact sequence,

$$\mathbf{0} \to K \to \operatorname{Ind} M \boxtimes L(i) \to \widetilde{f}_i M \to \mathbf{0} \tag{121}$$

and recall $\widetilde{f}_i M$ is the unique composition factor of $\operatorname{Ind} M \boxtimes L(i)$ such that $\varepsilon_i(\widetilde{f}_i M) = \varepsilon_i(M)+1$, and that for all composition factors N of K, $\varepsilon_i(N) \leq \varepsilon_i(M)$. By the exactness of induction there is a second short exact sequence

$$\mathbf{0} \to \operatorname{Ind} L(j) \boxtimes K \to \operatorname{Ind} L(j) \boxtimes M \boxtimes L(i) \to \operatorname{Ind} L(j) \boxtimes \widetilde{f}_i M \to \mathbf{0}, \tag{122}$$

and since $i \neq j$ the Shuffle Lemma tells us that for all composition factors N' of $\operatorname{Ind} L(j) \boxtimes K$, $\varepsilon_i(N') \leq \varepsilon_i(M)$. By the Shuffle Lemma and Frobenius reciprocity

$$\varepsilon_i(\widetilde{f}_j^\vee \widetilde{f}_i M) = \varepsilon_i(\widetilde{f}_i \widetilde{f}_j^\vee M) = \varepsilon_i(M) + 1. \tag{123}$$

Hence there can be no nonzero map

$$\operatorname{Ind} L(j) \boxtimes K \to \widetilde{f}_i \widetilde{f}_j^\vee M, \tag{124}$$

so that the submodule $\operatorname{Ind} L(j) \boxtimes K$ is contained in the kernel of β, as pictured in (125).

$$\begin{tikzcd}
 & \operatorname{Ind} L(j) \boxtimes \widetilde{f}_i M \arrow[r, two heads] & \widetilde{f}_j^\vee \widetilde{f}_i M \\
\operatorname{Ind} L(j) \boxtimes M \boxtimes L(i) \arrow[ur, two heads] \arrow[dr, two heads] \arrow[urr, bend left, two heads, "\alpha"] \arrow[drr, bend right, two heads, "\beta"'] & & \\
 & \operatorname{Ind} \widetilde{f}_j^\vee M \boxtimes L(i) \arrow[r, two heads] & \widetilde{f}_i \widetilde{f}_j^\vee M
\end{tikzcd} \tag{125}$$

Hence β induces a nonzero map (necessarily surjective)

$$\operatorname{Ind} L(j) \boxtimes \widetilde{f}_i M \twoheadrightarrow \widetilde{f}_i \widetilde{f}_j^{\vee} M. \tag{126}$$

Because $\operatorname{Ind} L(j) \boxtimes \widetilde{f}_i M$ has unique simple quotient $\widetilde{f}_j^{\vee} \widetilde{f}_i M$, then $\widetilde{f}_j^{\vee} \widetilde{f}_i M \cong \widetilde{f}_i \widetilde{f}_j^{\vee} M$. This proves 1a.

The three isomorphisms in 1b, 1c, and 1d all follow from 1a. For example, if $\widetilde{e}_j^{\vee} M$ is nonzero, then

$$\widetilde{f}_i M \cong \widetilde{f}_i \widetilde{f}_j^{\vee} \widetilde{e}_j^{\vee} M \cong \widetilde{f}_j^{\vee} \widetilde{f}_i \widetilde{e}_j^{\vee} M. \tag{127}$$

Applying $\widetilde{e}_j^{\vee}$ to both sides we get 1b. 1c and 1d follow similarly.

2. We prove 2a. Let $c = \varepsilon_i^{\vee}(M), m = \varepsilon_i(M)$.

   - Suppose $\operatorname{jump}_i(M) = 0$. Then also $\operatorname{jump}_i(\widetilde{f}_i M) = \operatorname{jump}_i(\widetilde{f}_i^{\vee} M) = 0$ by (117). Thus by Lemma 4.14
   $$\widetilde{f}_i^{\vee} \widetilde{f}_i M \cong \widetilde{f}_i \widetilde{f}_i M \cong \widetilde{f}_i \widetilde{f}_i^{\vee} M. \tag{128}$$
   - Suppose $\operatorname{jump}_i(M) = 1$. By Lemma 4.14 and Proposition 4.6, $\varepsilon_i(\widetilde{f}_i M) = m+1$ but $\varepsilon_i^{\vee}(\widetilde{f}_i M) = c$. While $\varepsilon_i(\widetilde{f}_i^{\vee} M) = m$ but $\varepsilon_i^{\vee}(\widetilde{f}_i^{\vee} M) = c+1$. Further by (117) $\operatorname{jump}_i(\widetilde{f}_i M) = \operatorname{jump}_i(\widetilde{f}_i^{\vee} M) = 0$. Hence $\varepsilon_i(\widetilde{f}_i^{\vee} \widetilde{f}_i M) = m+2$, $\varepsilon_i^{\vee}(\widetilde{f}_i^{\vee} \widetilde{f}_i M) = c+1$ whereas $\varepsilon_i(\widetilde{f}_i \widetilde{f}_i^{\vee} M) = m+1$, $\varepsilon_i^{\vee}(\widetilde{f}_i \widetilde{f}_i^{\vee} M) = c+2$. Thus the two modules cannot be isomorphic.
   - Suppose $\operatorname{jump}_i(M) \geq 2$. Then $\operatorname{jump}_i(\widetilde{f}_i M) = \operatorname{jump}_i(\widetilde{f}_i^{\vee} M) \geq 1$. We calculate
   $$\varepsilon_i(\widetilde{f}_i \widetilde{f}_i^{\vee} M) = m+1 = \varepsilon_i(\widetilde{f}_i^{\vee} \widetilde{f}_i M) \tag{129}$$
   $$\varepsilon_i^{\vee}(\widetilde{f}_i \widetilde{f}_i^{\vee} M) = c+1 = \varepsilon_i^{\vee}(\widetilde{f}_i^{\vee} \widetilde{f}_i M). \tag{130}$$

     We will show there is no nonzero map
   $$\operatorname{Ind} L(i) \boxtimes K \to \widetilde{f}_i \widetilde{f}_i^{\vee} M \tag{131}$$
     for any proper submodule $K \subseteq \operatorname{Ind} M \boxtimes L(i)$. Given we have a surjection
   $$\operatorname{Ind} L(i) \boxtimes M \boxtimes L(i) \twoheadrightarrow \widetilde{f}_i \widetilde{f}_i^{\vee} M \tag{132}$$
     this means we must have a nonzero map
   $$\operatorname{Ind} L(i) \boxtimes \widetilde{f}_i M \to \widetilde{f}_i \widetilde{f}_i^{\vee} M, \tag{133}$$
     which will prove the lemma as
   $$\widetilde{f}_i^{\vee} \widetilde{f}_i M = \operatorname{cosoc} \operatorname{Ind} L(i) \boxtimes \widetilde{f}_i M. \tag{134}$$

     First note there is no nonzero map
   $$\operatorname{Ind} L(i) \boxtimes \widetilde{f}_i^{\vee} M \to \widetilde{f}_i \widetilde{f}_i^{\vee} M \tag{135}$$
     as $\operatorname{cosoc}(\operatorname{Ind} L(i) \boxtimes \widetilde{f}_i^{\vee} M) = (\widetilde{f}_i^{\vee})^2 M$ and $\varepsilon_i^{\vee}((\widetilde{f}_i^{\vee})^2 M) = c+2 \neq c+1 = \varepsilon_i^{\vee}(\widetilde{f}_i \widetilde{f}_i^{\vee} M)$. Let D be any other composition factor of $\operatorname{Ind} M \boxtimes L(i)$ apart from $\widetilde{f}_i M$ or $\widetilde{f}_i^{\vee} M$

(recall the latter occur with multiplicity one as composition factors). Then by Proposition 4.6, $\varepsilon_i(D) \leq m$, $\varepsilon_i^\vee(D) \leq c$. If there were a nonzero map $\operatorname{Ind} L(i) \boxtimes D \to \widetilde{f}_i \widetilde{f}_i^\vee M$, it would imply $\widetilde{f}_i^\vee D \cong \widetilde{f}_i \widetilde{f}_i^\vee M$ and so $\varepsilon_i^\vee(\widetilde{f}_i^\vee D) = c + 1$ meaning $\varepsilon_i^\vee(D) = c$. Also $m + 1 = \varepsilon_i(\widetilde{f}_i^\vee D) \leq \varepsilon_i(D) + 1$ by the Shuffle Lemma, forcing $\varepsilon_i(D) = m$. By Lemma 4.14 this forces $0 = \mathrm{jump}_i(D)$ and $\widetilde{f}_i D \cong \widetilde{f}_i^\vee D \cong \widetilde{f}_i \widetilde{f}_i^\vee M$ from above, forcing $D \cong \widetilde{f}_i^\vee M$, which we already ruled out. Hence there must be a nonzero map

$$\operatorname{Ind} L(i) \boxtimes \widetilde{f}_i M \to \widetilde{f}_i \widetilde{f}_i^\vee M. \tag{136}$$

Now that we have established $\widetilde{f}_i^\vee \widetilde{f}_i M \cong \widetilde{f}_i \widetilde{f}_i^\vee M$ if and only if $\mathrm{jump}_i(M) \neq 1$, statements 2b and 2c follow directly from Proposition 4.6.6.

□

Remark 4.18. Because $\widetilde{e}_i$ and $\widetilde{f}_j^\vee$ commute for $i \neq j$, then $\varepsilon_i(\widetilde{f}_j^\vee M) = \varepsilon_i(M)$. An equivalent statement holds for $\widetilde{e}_i^\vee$, $\widetilde{f}_j$, and $\varepsilon_i^\vee$. When $\mathrm{jump}_i(M) \neq 0$, $\varepsilon_i(\widetilde{f}_i^\vee M) = \varepsilon_i(M)$.

Proposition 4.19. Let M be a simple $R(\nu)$-module. Let $i, j \in I$ and $a_{ij} = 0$. Then

1. $\widetilde{f}_i \widetilde{f}_j M \cong \widetilde{f}_j \widetilde{f}_i M$.
2. $\widetilde{e}_i \widetilde{e}_j M \cong \widetilde{e}_j \widetilde{e}_i M$.

Proof. The proof of part 1 is similar to the proof of Proposition 4.17.1a once we note that by formula (118) and Lemma 4.14,

$$\operatorname{Ind} L(i) \boxtimes L(j) \cong \operatorname{Ind} L(j) \boxtimes L(i). \tag{137}$$

Part 2 follows directly from part 1 by Proposition 4.6.6. Observe that Part 2 is true even in the case $\widetilde{e}_i \widetilde{e}_j M = \mathbf{0}$. □

5 Rep^Λ and the functor pr_Λ

All of the lemmas and propositions in this section about the interaction of the functors pr_Λ and Ind are adapted from [33].

For $\Lambda = \sum_{i \in I} \lambda_i \Lambda_i \in P^+$ define $\mathcal{I}_\nu^\Lambda$ to be the two-sided ideal of $R(\nu)$ generated by the elements $x_1^{\lambda_{i_1}} 1_{\underline{\mathbf{i}}}$ for all $\underline{\mathbf{i}} \in \mathrm{Seq}(\nu)$. When ν is clear from the context we write, $\mathcal{I}_\nu^\Lambda = \mathcal{I}^\Lambda$. The *cyclotomic KLR algebra of weight* Λ is then defined as

$$R^\Lambda = \bigoplus_{\nu \in Q^+} R^\Lambda(\nu) \quad \text{where} \quad R^\Lambda(\nu) := R(\nu)/\mathcal{I}_\nu^\Lambda. \tag{138}$$

The algebra $R^\Lambda(\nu)$ is finite dimensional, [4, 23]. The category of finite dimensional $R^\Lambda(\nu)$-modules is denoted $R^\Lambda(\nu)\text{-}\mathrm{mod}$ and the category of finite dimensional R^Λ-modules is denoted $R^\Lambda\text{-}\mathrm{mod}$. The category of finite dimensional R-modules on which $\mathcal{I}^\Lambda$ vanishes is denoted

$$\mathrm{Rep}^\Lambda.$$

While we can identify R^Λ - mod with Rep^Λ, we choose to work with Rep^Λ . We construct a right-exact functor, $\mathrm{pr}_\Lambda : R(\nu)\text{ - mod} \to R(\nu)\text{ - mod}$, via

$$\mathrm{pr}_\Lambda M := M/\mathcal{I}^\Lambda M \tag{139}$$

and extend it to $\mathrm{pr}_\Lambda : R\text{ - mod} \to R\text{ - mod}$. It is customary in the literature to interpret pr_Λ as being a functor from R - mod to R^Λ - mod, but in this paper it will be more convenient to consider it as a functor $R\text{ - mod} \to R\text{ - mod}$ or $R\text{ - mod} \to \mathrm{Rep}^\Lambda$. The reader may keep in mind that the image of pr_Λ consists of $R(\nu)$-modules which descend to $R^\Lambda(\nu)$-modules. Observe that in the opposite direction there is an exact functor $\mathrm{infl}_\Lambda : R^\Lambda(\nu)\text{ - mod} \to R(\nu)\text{ - mod}$, where $R(\nu)$ acts on $R^\Lambda(\nu)$-module M through the projection map $R(\nu) \twoheadrightarrow R^\Lambda(\nu)$.

Remark 5.1. If M is a $R(\nu)$-module and A is a simple module in Rep^Λ for $\Lambda \in P^+$, then since $\mathrm{pr}_\Lambda A \cong A$, the right exactness of pr_Λ implies that any surjection $M \twoheadrightarrow A$ gives a surjection $\mathrm{pr}_\Lambda M \twoheadrightarrow A$. Similarly, since there always exists a surjection $M \twoheadrightarrow \mathrm{pr}_\Lambda M$, given a surjection $\mathrm{pr}_\Lambda M \twoheadrightarrow A$ we immediately get a surjection $M \twoheadrightarrow A$. In such situations there is an equivalence between the two surjections $M \twoheadrightarrow A$ and $\mathrm{pr}_\Lambda M \twoheadrightarrow A$ which we will henceforth use freely.

If M is simple then either $\mathrm{pr}_\Lambda M = \mathbf{0}$ or $\mathrm{pr}_\Lambda M = M$. There is a useful criterion for determining the action of pr_Λ on simple $R(\nu)$-modules given by the following proposition.

Proposition 5.2. [23] Let $\Lambda = \sum_{i\in I} \lambda_i \Lambda_i \in P^+$, $\nu \in Q^+$, and let M be a simple $R(\nu)$-module. Then $\mathcal{I}^\Lambda M = \mathbf{0}$ if and only if $\mathrm{pr}_\Lambda M \cong M$ if and only if $\mathrm{pr}_\Lambda M \neq \mathbf{0}$ if and only if

$$\varepsilon_i^\vee(M) \leq \lambda_i$$

for all $i \in I$. When these conditions hold $M \in \mathrm{Rep}^\Lambda$, and we may identify M with $\mathrm{pr}_\Lambda M$ (or as an $R^\Lambda(\nu)$-module).

In this paper we will primarily consider $\Lambda = \Lambda_i$ in which case $\mathcal{I}_\nu^{\Lambda_i}$ is generated by $x_1 1_{ii_2\dots i_m}$ and $1_{ji_2\dots i_m}, j \neq i$ ranging over $\underline{\mathbf{i}} \in \mathrm{Seq}(\nu)$.

Notice that Proposition 5.2 immediately tells us that if p is a cyclotomic path of tail weight (Λ_j, Λ_i) then the modules $T_{p;k}$ belong to Rep^{Λ_i} for any $k \geq 0$. This is part of the motivation for the definition of cyclotomic path in Section 2.2.

For $\Lambda = \sum_{i\in I} \lambda_i \Lambda_i \in P^+$ and M an irreducible $R(\nu)$-module set

$$\varphi_j^\Lambda(M) = \lambda_j + \varepsilon_j(M) + \mathrm{wt}_j(M). \tag{140}$$

Notice that when $\Lambda = \Lambda_i$ this gives

$$\varphi_j^{\Lambda_i}(M) = \delta_{ij} + \varepsilon_j(M) + \mathrm{wt}_j(M). \tag{141}$$

Remark 5.3. By formula (141) if M is a simple module in Rep^{Λ_i} it follows that

$$\varphi_j^{\Lambda_i}(M) = \begin{cases} \delta_{ij} & \text{if } M = \mathbb{1}, \\ \mathrm{jump}_j(M) & \text{otherwise.} \end{cases} \tag{142}$$

Proposition 5.4. [23] Let M be a simple $R(\nu)$-module with $\mathrm{pr}_\Lambda M \neq \mathbf{0}$. Then

$$\varphi_j^\Lambda(M) = \max\{k \in \mathbb{Z} \mid \mathrm{pr}_\Lambda \widetilde{f}_j^k M \neq \mathbf{0}\}. \tag{143}$$

5.0.1 Module-theoretic model of $B(\Lambda)$

Let M be a simple $R(\nu)$-module. Set

$$\mathrm{wt}(M) = -\nu \quad \text{and} \quad \mathrm{wt}_i(M) = -\langle h_i, \nu\rangle. \tag{144}$$

Let $\mathrm{Irr}\, R$ be the set of isomorphism classes of simple R-modules and $\mathrm{Irr}\, R^\Lambda$ be the set of isomorphism classes of simple modules in Rep^Λ. In [23] it was shown that the tuple $(\mathrm{Irr}\, R, \varepsilon_i, \varphi_i, \widetilde{e}_i, \widetilde{f}_i, \mathrm{wt})$ defines a crystal isomorphic to $B(\infty)$ and $(\mathrm{Irr}\, R^\Lambda, \varepsilon_i, \varphi_i^\Lambda, \widetilde{e}_i, \widetilde{f}_i, \mathrm{wt})$ defines a crystal isomorphic to the highest weight crystal $B(\Lambda)$.

From property (117) of jump_i it is clear that if we apply $\widetilde{f}_i$ sufficiently many times to any simple module $M \in \mathrm{Rep}^\Lambda$, then eventually we will reach an n for which

$$\varepsilon_i^\vee(\widetilde{f}_i^n M) > \lambda_i \tag{145}$$

and so $\mathrm{pr}_\Lambda \widetilde{f}_i^n M = \mathbf{0}$. Proposition 5.4 says that φ_i^Λ measures this for simple modules in Rep^Λ. In fact it is true that $\mathrm{pr}_\Lambda M \neq \mathbf{0}$ if and only if $\varphi_i^\Lambda(M) \geq 0$ for all $i \in I$. Above we saw that the function φ_i^Λ is part of a crystal datum. With this is mind, we mimic the conventions usually used in the theory of crystals and define

$$\varphi^\Lambda(M) := \sum_{i\in I} \varphi_i^\Lambda(M)\Lambda_i, \tag{146}$$

and

$$\varepsilon(M) := \sum_{i\in I} \varepsilon_i(M)\Lambda_i, \qquad \varepsilon^\vee(M) := \sum_{i\in I} \varepsilon_i^\vee(M)\Lambda_i. \tag{147}$$

Lemma 5.5. Let M be an $R(\nu)$-module such that $\mathrm{pr}_\Lambda M \neq \mathbf{0}$. Let Q be a simple quotient of $\mathrm{pr}_\Lambda M$. Then $\mathrm{pr}_\Lambda Q = Q$.

Proof. This follows from the right exactness of pr_Λ. □

5.0.2 Interaction of pr_Λ and Ind

The following is a list of useful facts about the way that the functor pr_Λ interacts with induction.

Proposition 5.6. Let $\mu, \nu \in Q^+$, $\Lambda \in P^+$, $i, j \in I$, and let M be simple $R(\mu)$-module and N a simple $R(\nu)$-module.

(a) If $\mathrm{pr}_\Lambda M = \mathbf{0}$ then $\mathrm{pr}_\Lambda \mathrm{Ind}\, M \boxtimes N = \mathbf{0}$.

(b) If $\mathrm{pr}_\Lambda \mathrm{Ind}\, M \boxtimes L(i^c) = \mathbf{0}$ and $\varepsilon_i^\vee(N) \geq c$ then $\mathrm{pr}_\Lambda \mathrm{Ind}\, M \boxtimes N = \mathbf{0}$.

(c) If $c > \varphi_i^\Lambda(M)$ then $\mathrm{pr}_\Lambda \mathrm{Ind}\, M \boxtimes L(i^c) = \mathbf{0}$.

(d) If $\mathrm{pr}_\Lambda \mathrm{Ind}\, M \boxtimes N \neq \mathbf{0}$ then $N \in \mathrm{Rep}^{\varphi^\Lambda(M)}$.

(e) Let $\varphi = \varphi_i^\Lambda(M)$, then $\mathrm{pr}_\Lambda \mathrm{Ind}\, M \boxtimes L(i^\varphi) \cong \widetilde{f}_i^\varphi M$.

(f) If $\mathrm{pr}_\Lambda C = M$ then $\mathrm{pr}_\Lambda \mathrm{Ind}\, C \boxtimes N \cong \mathrm{pr}_\Lambda \mathrm{Ind}\, M \boxtimes N$.

(g) Suppose $a_{ij} < 0$, $\varepsilon(M) = \Lambda_i$, and $\mathrm{pr}_\Lambda \operatorname{Ind} \widetilde{e}_i M \boxtimes L(j) = \mathbf{0}$. Then $\mathrm{pr}_\Lambda \operatorname{Ind} M \boxtimes L(j) \cong \mathrm{pr}_\Lambda \widetilde{f}_j M$. In particular, if $\mathrm{pr}_\Lambda M \neq \mathbf{0}$ then $\varepsilon(\widetilde{f}_j M) = \Lambda_j$.

Proof. We write

$$\Lambda = \sum_{i \in I} \lambda_i \Lambda_i. \tag{148}$$

(a) If $\mathrm{pr}_\Lambda M = \mathbf{0}$, then by Proposition 5.2 there is some $i \in I$ such that $\varepsilon_i^\vee(M) > \lambda_i$. Suppose that $\mathrm{pr}_\Lambda \operatorname{Ind} M \boxtimes N \neq \mathbf{0}$. Then it has some simple quotient Q, and there are surjections

$$\operatorname{Ind} M \boxtimes N \twoheadrightarrow \mathrm{pr}_\Lambda \operatorname{Ind} M \boxtimes N \twoheadrightarrow Q. \tag{149}$$

By Lemma 5.5 $\mathrm{pr}_\Lambda Q = Q$. By Frobenius reciprocity $\operatorname{Res}^{\mu+\nu}_{\mu,\nu} Q$ has $M \boxtimes N$ as a $(R(\mu) \otimes R(\nu))$-submodule. But Remark 4.7 then implies $\varepsilon_i^\vee(Q) > \lambda_i$ so that $\mathrm{pr}_\Lambda Q = \mathbf{0}$, a contradiction.

(b) If $\varepsilon_i^\vee(N) \geq c$ then there is a surjection,

$$\operatorname{Ind} L(i^c) \boxtimes (\widetilde{e}_i^\vee)^c N \twoheadrightarrow N \tag{150}$$

and by the exactness of induction a surjection

$$\operatorname{Ind} M \boxtimes L(i^c) \boxtimes (\widetilde{e}_i^\vee)^c N \twoheadrightarrow \operatorname{Ind} M \boxtimes N. \tag{151}$$

If $\mathrm{pr}_\Lambda \operatorname{Ind} M \boxtimes L(i^c) = \mathbf{0}$, then by part (a) above and the right exactness of pr_Λ, $\mathrm{pr}_\Lambda \operatorname{Ind} M \boxtimes N = \mathbf{0}$.

(c) This follows from Proposition 5.4 and the fact that the induced module has unique simple quotient $\widetilde{f}_i^c M$; or see [23].

(d) We will show the contrapositive. Suppose $N \notin \mathrm{Rep}^{\varphi^\Lambda(M)}$. Then by Proposition 5.2, there is some $i \in I$ such that $\varepsilon_i^\vee(N) = c > \varphi_i^\Lambda(M)$. By part (c), $\mathrm{pr}_\Lambda \operatorname{Ind} M \boxtimes L(i^c) = \mathbf{0}$. Part (b) then implies that $\mathrm{pr}_\Lambda \operatorname{Ind} M \boxtimes N = \mathbf{0}$.

(e) Consider the exact sequence,

$$\mathbf{0} \to K \to \operatorname{Ind} M \boxtimes L(i^\varphi) \to \widetilde{f}_i^\varphi M \to \mathbf{0}. \tag{152}$$

$\widetilde{f}_i^\varphi M$ is the unique composition factor of $\operatorname{Ind} M \boxtimes L(i^\varphi)$ such that $\varepsilon_i(\widetilde{f}_i^\varphi M) = \varphi + \varepsilon_i(M)$, and $\varepsilon_i(D) < \varphi + \varepsilon_i(M)$ for all composition factors D of K. All composition factors D of K have the same weight as $\widetilde{f}_i^\varphi M$. By (140) and Proposition 4.6, $\varphi_i^\Lambda(D) = \lambda_i + \varepsilon_i(D) + \mathrm{wt}_i(D) < \lambda_i + \varepsilon_i(\widetilde{f}_i^\varphi M) + \mathrm{wt}_i(\widetilde{f}_i^\varphi M) = \varphi_i^\Lambda(\widetilde{f}_i^\varphi M) = 0$. In particular this shows $\mathrm{pr}_\Lambda K = \mathbf{0}$ so by the right exactness of pr_Λ we get (e).

(f) Consider the diagram in (153),

$$\begin{tikzcd}
& & \mathbf{0} \arrow[d] & & \\
& & \operatorname{Ind}\mathcal{I}^{\Lambda}C \boxtimes N \arrow[d, "\alpha"'] \arrow[dr, "\beta\circ\alpha"] & & \\
\mathbf{0} \arrow[r] & \mathcal{I}^{\Lambda}(\operatorname{Ind}C \boxtimes N) \arrow[r] & \operatorname{Ind}C \boxtimes N \arrow[r, "\beta"] \arrow[d, "\gamma"'] & \mathrm{pr}_{\Lambda}(\operatorname{Ind}C \boxtimes N) \arrow[r] & \mathbf{0} \\
& & \operatorname{Ind}M \boxtimes N \arrow[ur, dotted, "g"'] \arrow[d] & & \\
& & \mathbf{0} & &
\end{tikzcd} \tag{153}$$

where the horizontal and vertical sequences are exact. Recall that $\mathcal{I}^{\Lambda}_{\mu}$ in $R(\mu)$ is generated by the set $\{x_1^{\lambda_{i_1}} 1_{\underline{\mathbf{i}}}\}_{\underline{\mathbf{i}}\in\mathrm{Seq}(\mu)}$ where $\underline{\mathbf{i}} = i_1 i_2 \ldots i_m$ and $|\mu| = m$. Under the embedding

$$R(\mu) \hookrightarrow R(\mu) \otimes R(\nu) \hookrightarrow R(\mu+\nu), \tag{154}$$

this set maps to the set

$$\Big\{ \sum_{\underline{\mathbf{j}}\in\mathrm{Seq}(\nu)} x_1^{\lambda_{i_1}} 1_{\underline{\mathbf{ij}}} \Big\}_{\underline{\mathbf{i}}\in\mathrm{Seq}(\mu)} \tag{155}$$

in $R(\mu+\nu)$. This set is contained in the ideal generated by $\{x_1^{\lambda_{i_1}} 1_{\underline{\mathbf{k}}}\}_{\underline{\mathbf{k}}\in\mathrm{Seq}(\mu+\nu)}$ which generates $\mathcal{I}^{\Lambda}_{\mu+\nu}$. It follows that

$$R(\mu+\nu)\mathcal{I}^{\Lambda}_{\mu} \subseteq \mathcal{I}^{\Lambda}_{\mu+\nu}, \tag{156}$$

and hence

$$\operatorname{Ind}\mathcal{I}^{\Lambda}_{\mu}C \boxtimes N \subseteq \mathcal{I}^{\Lambda}_{\mu+\nu}(\operatorname{Ind}C \boxtimes N). \tag{157}$$

This tells us that the composition $\beta\circ\alpha$ from the diagram in (153) is zero, so there exists a surjective homomorphism $g : \operatorname{Ind}M \boxtimes N \to \mathrm{pr}_{\Lambda}\operatorname{Ind}C \boxtimes N$. Applying pr_{Λ} to the diagram in (153), and sending maps γ, β, and g to $\tilde{\gamma}$, $\tilde{\beta}$, and $\tilde{g}$ respectively, right exactness yields $\tilde{\gamma}$, $\tilde{\beta}$, and $\tilde{g}$ are surjections as shown in diagram (158). It follows from considerations of dimension and that $\mathrm{pr}_{\Lambda} C = M$ that $\tilde{g}$ must be an isomorphism.

$$\begin{tikzcd}
\mathrm{pr}_{\Lambda}(\operatorname{Ind}C \boxtimes N) \arrow[r, "\tilde{\beta}"] \arrow[d, "\tilde{\gamma}"'] & \mathrm{pr}_{\Lambda}(\operatorname{Ind}C \boxtimes N) \arrow[r] & \mathbf{0} \\
\mathrm{pr}_{\Lambda}(\operatorname{Ind}M \boxtimes N) \arrow[ur, "\tilde{g}"'] \arrow[d] & & \\
\mathbf{0} & &
\end{tikzcd} \tag{158}$$

(g) First, $\varepsilon(M) = \Lambda_i$ implies $e_i M \cong \widetilde{e}_i M \neq \mathbf{0}$. Further, if $\mathrm{pr}_\Lambda \widetilde{e}_i M = \mathbf{0}$ then $\mathrm{pr}_\Lambda M = \mathbf{0}$ so the conclusion follows as all modules in question are $\mathbf{0}$. If $\mathrm{pr}_\Lambda \widetilde{e}_i M \neq \mathbf{0}$ the hypotheses imply that $\mathrm{Ind}\, \widetilde{e}_i M \boxtimes L(j)$ is irreducible by Lemma 4.14.

Suppose

$$\mathbf{0} \to K \to \mathrm{Ind}\, M \boxtimes L(j) \to \mathrm{pr}_\Lambda \mathrm{Ind}\, M \boxtimes L(j) \to \mathbf{0} \tag{159}$$

is exact. By the exactness of e_i then

$$\mathbf{0} \to e_i K \to e_i\Big(\mathrm{Ind}\, M \boxtimes L(j)\Big) \xrightarrow{\varphi} e_i\Big(\mathrm{pr}_\Lambda \mathrm{Ind}\, M \boxtimes L(j)\Big) \to \mathbf{0} \tag{160}$$

is also exact. By Proposition 4.9, $e_i(\mathrm{Ind}\, M \boxtimes L(j)) \cong \mathrm{Ind}\, e_i M \boxtimes L(j) \cong \mathrm{Ind}\, \widetilde{e}_i M \boxtimes L(j)$. But $\mathrm{pr}_\Lambda \mathrm{Ind}\, \widetilde{e}_i M \boxtimes L(j) = \mathbf{0}$ by hypothesis so the surjection φ then forces $e_i(\mathrm{pr}_\Lambda \mathrm{Ind}\, M \boxtimes L(j)) = \mathbf{0}$. In particular $e_i(\mathrm{pr}_\Lambda \widetilde{f}_j M) = \mathbf{0}$ and so $\varepsilon_i(\widetilde{f}_j M) = 0$.

Note if D is any composition factor of $\mathrm{Ind}\, M \boxtimes L(j)$ other than $\widetilde{f}_j M$ then $\varepsilon_j(D) = 0$ by Proposition 4.6.2 and the fact that $\varepsilon_j(M) = 0$. Then $\varepsilon(M) = \Lambda_i$ and the Shuffle Lemma forces $e_i D \neq \mathbf{0}$. Hence D cannot be a composition factor of $\mathrm{pr}_\Lambda \mathrm{Ind}\, M \boxtimes L(j)$. In particular $\mathrm{pr}_\Lambda \mathrm{Ind}\, M \boxtimes L(j) \cong \mathrm{pr}_\Lambda \widetilde{f}_j M$.

If $\mathrm{pr}_\Lambda \widetilde{f}_j M \neq \mathbf{0}$ then $\mathrm{pr}_\Lambda \widetilde{f}_j M = \widetilde{f}_j M$ because $\widetilde{f}_j M$ is simple and as $\varepsilon_j(\widetilde{f}_j M) = 1$, $\varepsilon_i(\widetilde{f}_j M) = 0$, and $\varepsilon(M) = \Lambda_i$ we must have $\varepsilon(\widetilde{f}_j M) = \Lambda_j$.

□

6 The family of modules $T_{p;k}$

In this section we describe certain modules, motivated by their crystal-theoretic description in [32], see Proposition 6.4 below. For each of the classical affine types listed in Section 2.1 we specify a set of elements $\{\gamma_{p;k}\}$ of Q^+ corresponding to paths in $B^{1,1}$. For path $p : \{0, 1, \ldots, k-1\} \to I$ of length k define

$$\gamma_{p;k} := \alpha_{p(0)} + \alpha_{p(1)} + \cdots + \alpha_{p(k-1)}. \tag{161}$$

Observe that $|\gamma_{p;k}| = k$. We also define $\underline{\tau_{\mathbf{p;k}}} \in \mathrm{Seq}(\gamma_{p;k})$ to be

$$\underline{\tau_{\mathbf{p;k}}} = (p(0), p(1), \ldots, p(k-1)). \tag{162}$$

Let $\eta_{j_1}^{j_2}(\underline{\tau_{\mathbf{p;k}}}) \subseteq \{1, 2, \ldots, k\}$ be the collection of indices for adjacent $j_1,\ j_2 \in I$ in the sequence $\underline{\tau_{\mathbf{p;k}}}$,

$$\eta_{j_1}^{j_2}(\underline{\tau_{\mathbf{p;k}}}) := \{\ t \mid p(t-1) = j_1,\ p(t) = j_2,\ 1 \le t \le k-1\}. \tag{163}$$

Let $|\eta_{j_1}^{j_2}(\underline{\tau_{\mathbf{p;k}}})|$ denote the cardinality of this set.

Example 6.1. If p is a type $A_6^{(2)}$ path such that

$$\underline{\tau_{\mathbf{p;18}}} = (0, 1, 2, 3, 2, 1, 0, 0, 1, 2, 3, 2, 1, 0, 0, 1, 2, 3), \tag{164}$$

then $\eta_0^0(\underline{\tau_{\mathbf{p;18}}}) = \{7, 14\}$.

To each path p of length k, associate an $R(\gamma_{p;k})$-module denoted by $T_{p;k}$. In all types $T_{p;0} = \mathbb{1}$, the unique simple $R(0)$-module which we will refer to as the *unit module.*

- $(A_\ell^{(1)})$: $T_{p;k}$ is a 1-dimensional simple module with character

$$\mathrm{Char}(T_{p;k}) = [\underline{\tau_{\mathbf{p};\mathbf{k}}}] = [p(0), p(1), \ldots, p(k-1)]. \tag{165}$$

If v spans $T_{p;k}$, then the generators of $R(\gamma_{p;k})$ act by

$$1_{\underline{\mathbf{i}}}v = \delta_{\underline{\mathbf{i}},\underline{\tau_{\mathbf{p};\mathbf{k}}}}v, \qquad x_r 1_{\underline{\mathbf{i}}}v = 0, \qquad \psi_r 1_{\underline{\mathbf{i}}}v = 0. \tag{166}$$

- $(C_\ell^{(1)})$: In this type $T_{p;k}$ is also 1-dimensional, again with character $\mathrm{Char}(T_{p;k}) = [\underline{\tau_{\mathbf{p};\mathbf{k}}}]$. The action of the generators of $R(\gamma_{p;k})$ is the same as in type $A_\ell^{(1)}$.

- $(A_{2\ell}^{(2)})$: Let $d = |\eta_0^0(\underline{\tau_{\mathbf{p};\mathbf{k}}})|$. Then each $T_{p;k}$ has ungraded dimension 2^d and character

$$\mathrm{Char}(T_{p;k}) = q_0^d[2]_0^d[\underline{\tau_{\mathbf{p};\mathbf{k}}}]. \tag{167}$$

If

$$\eta_0^0(\underline{\tau_{\mathbf{p};\mathbf{k}}}) = \{t_1, \ldots, t_d\} \tag{168}$$

then $T_{p;k}$ has homogeneous basis

$$\left\{ v_\beta \;\middle|\; \beta = (\beta_1, \ldots, \beta_d) \in \{0,1\}^d \right\}. \tag{169}$$

The basis element v_β belongs to the degree $2(\beta_1 + \cdots + \beta_d)$ component of $T_{p;k}$. The generators of $R(\gamma_{p;k})$ act on v_β by

$$1_{\underline{\mathbf{i}}}v_\beta = \delta_{\underline{\mathbf{i}},\underline{\tau_{\mathbf{p};\mathbf{k}}}}v_\beta, \tag{170}$$

$$x_r v_\beta = x_r 1_{\underline{\tau_{\mathbf{p};\mathbf{k}}}} v_\beta = \begin{cases} v_{\beta+\delta_r} & \text{if } r \in \eta_0^0(\underline{\tau_{\mathbf{p};\mathbf{k}}}),\ \beta_r = 0 \\ -v_{\beta+\delta_r} & \text{if } r-1 \in \eta_0^0(\underline{\tau_{\mathbf{p};\mathbf{k}}}),\ \beta_r = 0 \\ 0 & \text{otherwise} \end{cases} \tag{171}$$

(here δ_r is the element of $\{0,1\}^d$ with only rth coordinate non-zero). Finally,

$$\psi_r v_\beta = \psi_r 1_{\underline{\tau_{\mathbf{p};\mathbf{k}}}} v_\beta = \begin{cases} v_{\beta-\delta_r} & \text{if } r \in \eta_0^0(\underline{\tau_{\mathbf{p};\mathbf{k}}}),\ \beta_r = 1 \\ 0 & \text{otherwise.} \end{cases} \tag{172}$$

- $(A_{2\ell}^{(2)\dagger})$: Let $d = |\eta_\ell^\ell(\underline{\tau_{\mathbf{p};\mathbf{k}}})|$. Then $T_{p;k}$ has ungraded dimension 2^d and character

$$\mathrm{Char}(T_{p;k}) = q_\ell^d[2]_\ell^d[\underline{\tau_{\mathbf{p};\mathbf{k}}}]. \tag{173}$$

There is a basis for $T_{p;k}$ whose description is similar to the description of the basis $\{v_\beta\}$ for type $A_{2\ell}^{(2)}$, the only difference being that we replace $\eta_0^0(\underline{\tau_{\mathbf{p};\mathbf{k}}})$ with $\eta_\ell^\ell(\underline{\tau_{\mathbf{p};\mathbf{k}}})$ everywhere.

- $(D^{(2)}_{\ell+1})$: Let $d = |\eta^0_0(\underline{\tau_{\mathbf{p};\mathbf{k}}})| + |\eta^\ell_\ell(\underline{\tau_{\mathbf{p};\mathbf{k}}})|$. Here each $T_{p;k}$ has ungraded dimension 2^d and character
$$\mathrm{Char}(T_{p;k}) = q_0^d[2]_0^d[\underline{\tau_{\mathbf{p};\mathbf{k}}}] \tag{174}$$
(note that in this type $[2]_0 = [2]_\ell$ since $(\alpha_0, \alpha_0) = (\alpha_\ell, \alpha_\ell) = 2$, so we may write formula (174) in terms of $[2]_0$ instead of both $[2]_0$ and $[2]_\ell$). It has a homogeneous basis analogous to that for types $A^{(2)}_{2\ell}$ and $A^{(2)\dagger}_{2\ell}$ which we again call $\{v_\beta\}$. This time however we replace $\eta^0_0(\underline{\tau_{\mathbf{p};\mathbf{k}}})$ from the description for type $A^{(2)}_{2\ell}$ with $\eta^0_0(\underline{\tau_{\mathbf{p};\mathbf{k}}}) \cup \eta^\ell_\ell(\underline{\tau_{\mathbf{p};\mathbf{k}}})$, and v_β belongs to the degree $2(\beta_1 + \cdots + \beta_d)$ component of $T_{p;k}$.

- $(A^{(2)}_{2\ell-1})$ Let $d = |\eta^1_0(\underline{\tau_{\mathbf{p};\mathbf{k}}}) \cup \eta^0_1(\underline{\tau_{\mathbf{p};\mathbf{k}}})|$. Here $T_{p;k}$ has graded dimension 2^d. If
$$\eta^1_0(\underline{\tau_{\mathbf{p};\mathbf{k}}}) \cup \eta^0_1(\underline{\tau_{\mathbf{p};\mathbf{k}}}) = \{t_1, \dots, t_d\} \tag{175}$$
then $T_{p;k}$ is concentrated in a single degree and has character
$$\mathrm{Char}(T_{p;k}) = \sum_{(\beta_1,\dots,\beta_d)\in\{0,1\}^d} \left[s_{t_1}^{\beta_1} s_{t_2}^{\beta_2} \dots s_{t_d}^{\beta_d} \underline{\tau_{\mathbf{p};\mathbf{k}}}\right]. \tag{176}$$
Here s_i is the adjacent transposition from $\mathcal{S}_k$ that interchanges i and $i+1$. $\mathcal{S}_k$ acts on $\mathrm{Seq}(\gamma_{p;k})$ as defined in (44). Let v be a nonzero vector from $1_{\underline{\tau_{\mathbf{p};\mathbf{k}}}} T_{p;k}$. Then a homogeneous weight basis for $T_{p;k}$ is given by
$$\{\psi_{t_1}^{\beta_1} \psi_{t_2}^{\beta_2} \dots \psi_{t_d}^{\beta_d} v \mid (\beta_1, \dots, \beta_d) \in \{0,1\}^d\}. \tag{177}$$
If $\beta = (\beta_1, \dots, \beta_d) \in \{0,1\}^d$ then write $v_\beta = \psi_{t_1}^{\beta_1} \psi_{t_2}^{\beta_2} \dots \psi_{t_d}^{\beta_d} v$. Also write
$$\underline{\mathbf{i}_\beta} = s_{t_1}^{\beta_1} s_{t_2}^{\beta_2} \dots s_{t_d}^{\beta_d} \underline{\tau_{\mathbf{p};\mathbf{k}}}. \tag{178}$$
Notice that the $\underline{\mathbf{i}_\beta}$ are distinct and $\dim(1_{\underline{\mathbf{i}_\beta}} T) = 1$. Then the generators of $R(\gamma_{p;k})$ act by,
$$1_{\underline{\mathbf{i}}} v_\beta = \delta_{\underline{\mathbf{i}}, \underline{\mathbf{i}_\beta}} v_\beta. \tag{179}$$
$$x_r v_\beta = x_r 1_{\underline{\mathbf{i}_\beta}} v_\beta = 0, \quad 1 \le r \le k, \tag{180}$$
$$\psi_r v_\beta = \psi_r 1_{\underline{\mathbf{i}_\beta}} v_\beta = \begin{cases} v_{\beta + \delta_r \bmod 2} & \text{if } r \in \eta^1_0(\underline{\tau_{\mathbf{p};\mathbf{k}}}) \cup \eta^0_1(\underline{\tau_{\mathbf{p};\mathbf{k}}}), \\ 0, & \text{otherwise.} \end{cases} \tag{181}$$

- $(B^{(1)}_\ell)$ For this type write $d_1 = |\eta^1_0(\underline{\tau_{\mathbf{p};\mathbf{k}}}) \cup \eta^0_1(\underline{\tau_{\mathbf{p};\mathbf{k}}})|$ and $d_2 = |\eta^\ell_\ell(\underline{\tau_{\mathbf{p};\mathbf{k}}})|$. Here $T_{p;k}$ has ungraded dimension $2^{d_1+d_2}$. If
$$\eta^1_0(\underline{\tau_{\mathbf{p};\mathbf{k}}}) \cup \eta^0_1(\underline{\tau_{\mathbf{p};\mathbf{k}}}) = \{t_1, \dots, t_{d_1}\} \tag{182}$$
$$\eta^\ell_\ell(\underline{\tau_{\mathbf{p};\mathbf{k}}}) = \{t'_1, \dots, t'_{d_2}\}. \tag{183}$$
Then $T_{p;k}$ has character
$$\mathrm{Char}(T_{p;k}) = \sum_{(\beta_1,\dots,\beta_{d_1})\in\{0,1\}^{d_1}} q_\ell^{d_2}[2]_\ell^{d_2} \left[s_{t_1}^{\beta_1} s_{t_2}^{\beta_2} \dots s_{t_{d_1}}^{\beta_{d_1}} \underline{\tau_{\mathbf{p};\mathbf{k}}}\right]. \tag{184}$$
We can describe a homogeneous weight basis $\{v_\beta\}$ for $T_{p;k}$ and action by the generators of $R(\gamma_{p;k})$ in this type by mixing our descriptions for type $A^{(2)\dagger}_{2\ell}$ with type $A^{(2)}_{2\ell-1}$.

- $(D_\ell^{(1)})$: Let $d = |\eta_0^1(\underline{\tau_{\mathbf{p};\mathbf{k}}}) \cup \eta_1^0(\underline{\tau_{\mathbf{p};\mathbf{k}}}) \cup \eta_{\ell-1}^{\ell}(\underline{\tau_{\mathbf{p};\mathbf{k}}}) \cup \eta_\ell^{\ell-1}(\underline{\tau_{\mathbf{p};\mathbf{k}}})|$. Here $T_{p;k}$ has graded dimension 2^d. If
$$\eta_0^1(\underline{\tau_{\mathbf{p};\mathbf{k}}}) \cup \eta_1^0(\underline{\tau_{\mathbf{p};\mathbf{k}}}) \cup \eta_{\ell-1}^{\ell}(\underline{\tau_{\mathbf{p};\mathbf{k}}}) \cup \eta_\ell^{\ell-1}(\underline{\tau_{\mathbf{p};\mathbf{k}}}) = \{t_1, \dots, t_d\} \tag{185}$$
then $T_{p;k}$ is concentrated in a single degree which we can take to be zero and has character
$$\mathrm{Char}(T_{p;k}) = \sum_{(\beta_1,\dots,\beta_d)\in\{0,1\}^d} \left[s_{t_1}^{\beta_1} s_{t_2}^{\beta_2} \dots s_{t_d}^{\beta_d} \underline{\tau_{\mathbf{p};\mathbf{k}}}\right]. \tag{186}$$
We can describe a homogeneous weight basis $\{v_\beta\}$ for $T_{p;k}$ and action on it by the generators for $R(\gamma_{p;k})$ in this type in the same manner as we did for type $A_{2\ell-1}^{(2)}$ by replacing every instance of $\eta_0^1(\underline{\tau_{\mathbf{p};\mathbf{k}}}) \cup \eta_1^0(\underline{\tau_{\mathbf{p};\mathbf{k}}})$ with $\eta_0^1(\underline{\tau_{\mathbf{p};\mathbf{k}}}) \cup \eta_1^0(\underline{\tau_{\mathbf{p};\mathbf{k}}}) \cup \eta_{\ell-1}^{\ell}(\underline{\tau_{\mathbf{p};\mathbf{k}}}) \cup \eta_\ell^{\ell-1}(\underline{\tau_{\mathbf{p};\mathbf{k}}})$.

Remark 6.2. In the above description of characters, we could have removed global factors such as q_i^d, as we only care about the simples up to overall grading shift. Further, in those cases above we in fact have $q_i = q$.

Showing that all these descriptions actually define modules involves checking that the actions agree with the KLR relations. For the $T_{p;k}$ above which are 1-dimensional, it is obvious that $T_{p;k}$ is simple. Showing that $T_{p;k}$ is simple in each of the other types is an easy exercise.

By [23], a simple R-module corresponds to a node in a highest weight crystal and the nodes corresponding to the $T_{p;k}$ were precisely those studied in [32]. See Proposition 6.4 below.

In type $A_\ell^{(1)}$ we could have also chosen to work with $B^{\ell,1}$. As noted in Remark 2.1 in type $A_\ell^{(1)}$, $B^{\ell,1}$ is similar to $B^{1,1}$ but with all arrows reversed. If p is a path in $B^{\ell,1}$ then as above define
$$\gamma_{p;k}^- := \alpha_{p(0)} + \cdots + \alpha_{p(k-1)} \qquad \underline{\tau_{\mathbf{p};\mathbf{k}}^-} = (p(0), \dots, p(k-1)), \tag{187}$$
and let $S_{p;k}$ denote the simple 1-dimensional $R(\gamma_{p;k}^-)$-module with character
$$\mathrm{Char}(S_{p;k}) = [\underline{\tau_{\mathbf{p};\mathbf{k}}^-}]. \tag{188}$$
In the Hecke algebra or symmetric group case, $T_{p;k}$ are the trivial modules and $S_{p;k}$ are the sign modules (see [33]).

Remark 6.3. In fact, given any 1-dimensional $R(\nu)$-module M in type $A_\ell^{(1)}$, there is some path p in either $B^{1,1}$ or $B^{\ell,1}$ such that $\nu = \gamma_{p;k}$ and $M \cong T_{p;k}$.

Returning to our family of modules $T_{p;k}$ from Section 4.1 we note that the path p that parametrizes $T_{p;k}$ also determines the sequence of $\widetilde{f}_i$ that we have to apply to $\mathbb{1}$ to construct to $T_{p;k}$.

Proposition 6.4. For all types X_ℓ and for a fixed path p of length k in $B^{1,1}$
$$\widetilde{f}_{p(k-1)} \widetilde{f}_{p(k-2)} \cdots \widetilde{f}_{p(0)}\, \mathbb{1} \cong T_{p;k}. \tag{189}$$

Proof. Recall that

$$[p(0), p(1), \dots, p(k-1)] \tag{190}$$

is in the support of $T_{p;k}$ For $k \geq 1$, let $i = p(k-1)$. Inspection of $[\underline{\mathbf{i}}] \in \operatorname{supp}(T_{p;k})$ implies that $\varepsilon := \varepsilon_i(T_{p;k}) \in \{1, 2\}$ and in particular is nonzero. Also by inspection $\operatorname{Char}(\widetilde{e}_i^{\varepsilon_i} T_{p;k}) = \operatorname{Char}(e_i^{(\varepsilon)} T_{p;k})$ agrees with $\operatorname{Char}(T_{p;k-\varepsilon})$ (up to overall grading shift). Hence by Remark 4.2, $\widetilde{e}_i^{\varepsilon} T_{p;k} \cong T_{p;k-\varepsilon}$. Proposition 4.6.6 then implies $T_{p;k} \cong \widetilde{f}_i^{\varepsilon} T_{p;k-\varepsilon}$. The identity (189) then follows by induction.

□

Note that this proposition also implies that if $j \in \operatorname{ext}_p^+$ then $\widetilde{f}_j T_{p;k} \cong T_{p\star\pi(j);k+1}$.

We will frequently need to compute $\operatorname{jump}_j(T_{p;k})$ and $\varphi_j^{\Lambda_{p(0)}}(T_{p;k})$ for a path p in $B^{1,1}$. In our main theorems of Section 7, we will only consider paths p such that $T_{p;k} \in \operatorname{Rep}^{\Lambda_{p(0)}}$. In this case, Remarks 6.6 and 5.3 apply, so we can in fact just consider jump. However, for completeness, we give data for other p. Observe the right/left symmetry of jump_j as in Table 2 that is broken for $\varphi_j^{\Lambda_{p(0)}}(T_{p;k})$ in general.

We introduce special notation,

$$\varphi_j(p;k) := \varphi_j^{\Lambda_{p(0)}}(T_{p;k}), \qquad \varphi(p;k) = \varphi^{\Lambda_{p(0)}}(T_{p;k}). \tag{191}$$

When $k = 0$, the unit module $T_{p;0} = \mathbb{1}$ has

$$\operatorname{jump}_j(\mathbb{1}) = 0 \quad \text{and} \quad \varphi_j^{\Lambda_i}(\mathbb{1}) = \delta_{ij}.$$

When $k = 1$,

$$\begin{gathered}\operatorname{jump}_j(L(j)) = 0 = \varphi_j^{\Lambda_j}(L(j)) \quad \text{and} \\ \operatorname{jump}_j(L(i)) = -\langle h_j, \alpha_i \rangle = -a_{ji} = \varphi_j^{\Lambda_i}(L(i)).\end{gathered}$$

For $k \geq 2$, we return to the classification of elements of I summarized in Table 1. Tables 2 and 3 give the values of $\operatorname{jump}_j(T_{p;k})$ and $\varphi_j(p;k)$ for $k \geq 2$, according to the class that j belongs to. In these tables we use the notation that for $j \in I$

$$\delta_{j,\operatorname{ext}_p^-} = \begin{cases} 1 & \text{if } j \in \operatorname{ext}_p^- \\ 0 & \text{otherwise} \end{cases} \qquad \delta_{j,\operatorname{ext}_p^+} = \begin{cases} 1 & \text{if } j \in \operatorname{ext}_p^+ \\ 0 & \text{otherwise} \end{cases} \tag{192}$$

For $k > 1$, we define $\varphi_j^-(p;k)$ and $\varphi_j^+(p;k)$ as in Table 4, so that

$$\varphi_j(p;k) = \varphi_j^-(p;k) + \varphi_j^+(p;k). \tag{193}$$

Note $\varphi_j^-(p;k)$ is "controlled" by elements at the left end of the sequence $\underline{\tau_{\mathbf{p};\mathbf{k}}}$ and $\varphi_j^+(p;k)$ is controlled by elements at the right end of $\underline{\tau_{\mathbf{p};\mathbf{k}}}$. We also define,

$$\varphi^-(p;k) := \sum_{i \in I} \varphi_i^-(p;k)\Lambda_i \tag{194}$$

$$\varphi^+(p;k) := \sum_{i \in I} \varphi_i^+(p;k)\Lambda_i \tag{195}$$

so that

$$\varphi(p;k) = \varphi^-(p;k) + \varphi^+(p;k). \tag{196}$$

Class	$\mathrm{jump}_j(T_{p;k})$
class $\mathcal{A}$	$\delta_{j,\mathrm{ext}_p^-} + \delta_{j,\mathrm{ext}_p^+}$
class $\mathcal{B}$	$\delta_{j,\mathrm{ext}_p^-}(2 - \delta_{j,p(0)}) + \delta_{j,\mathrm{ext}_p^+}(2 - \delta_{j,p(k-1)})$
class $\mathcal{D}$	$\delta_{j,\mathrm{ext}_p^-} + \delta_{j,\mathrm{ext}_p^+}$

Table 2: Values of $\mathrm{jump}_j(T_{p;k})$, $k > 1$, by class of $j \in I$.

Class	$\varphi_j(p;k)$
class $\mathcal{A}$	$\delta_{j,\mathrm{ext}_p^-} + \delta_{j,\mathrm{ext}_p^+}$
class $\mathcal{B}$	$2\delta_{j,\mathrm{ext}_p^-} - \delta_{j,p(0)} + \delta_{j,\mathrm{ext}_p^+}(2 - \delta_{j,p(k-1)})$
class $\mathcal{D}$	$\delta_{j,\mathrm{ext}_p^-} + (\delta_{j,\mathrm{ext}_p^-} - 1)\delta_{j',p(0)} + \delta_{j,\mathrm{ext}_p^+}$

Table 3: Values of $\varphi_j(p;k)$, $k > 1$, by class of $j \in I$. When j is Class $\mathcal{D}$, (j, j') is a Class $\mathcal{D}$ pair.

Class	$\varphi_j^-(p;k)$	$\varphi_j^+(p;k)$
class $\mathcal{A}$	$\delta_{j,\mathrm{ext}_p^-}$	$\delta_{j,\mathrm{ext}_p^+}$
class $\mathcal{B}$	$2\delta_{j,\mathrm{ext}_p^-} - \delta_{j,p(0)}$	$\delta_{j,\mathrm{ext}_p^+}(2 - \delta_{j,p(k-1)})$
class $\mathcal{D}$	$\delta_{j,\mathrm{ext}_p^-} + (\delta_{j,\mathrm{ext}_p^-} - 1)\delta_{j',p(0)}$	$\delta_{j,\mathrm{ext}_p^+}$

Table 4: Values of $\varphi_j^-(p;k)$ and $\varphi_j^+(p;k)$, $k > 1$, by class of $j \in I$. When j is Class $\mathcal{D}$, (j, j') is a Class $\mathcal{D}$ pair.

Remark 6.5. There is a more visual interpretation of $\varphi_j(p;k)$, $\varphi_j^-(p;k)$, and $\varphi_j^+(p;k)$ in terms of the path p in $B^{1,1}$ when $k > 1$.

- The value $\varphi_j(p;k)$ is the maximum number of j-arrows that extend path p from both its tail and head.
- When $\varepsilon^\vee(T_{p;k}) = \Lambda_{p(0)}$, the value $\varphi_j^-(p;k)$ is the maximum number of j-arrows that extend the tail of path p,
$$\varphi_j^-(p;k) = \max\{k \in \mathbb{Z}_{\geq 0} \mid (\pi(j))^k \star p \neq \mathbf{0}\ \}. \tag{197}$$
- The value $\varphi_j^+(p;k)$ is the maximum number of j-arrows that extend the head of path p,
$$\varphi_j^+(p;k) = \max\{k \in \mathbb{Z}_{\geq 0} \mid p \star (\pi(j))^k \neq \mathbf{0}\ \}. \tag{198}$$

Observe that under the above hypotheses $\varphi_j^-(p;k)$ and $\varphi_j^+(p;k)$ take values in $\{0,1,2\}$. This agrees with the fact that for all types there is no $j \in I$ such that more than 2 j-arrows appear consecutively in $B^{1,1}$. (More generally, we can also have $\varphi_j^+(p;k) = -1$ when $T_{p;k} \notin \mathrm{Rep}^{\Lambda_{p(0)}}$).

When $T_{p;k} \notin \mathrm{Rep}^{\Lambda_{p(0)}}$ then $T_{p;k} \in \mathrm{Rep}^{\Lambda_{p(0)}+\Lambda_{p(1)}}$. Since $\langle h_{p(1)}, \Lambda_{p(0)}\rangle = \varepsilon^\vee_{p(1)}(T_{p;k}) - 1$ in this case, by (141) and (118) $\varphi_{p(1)}^{\Lambda_{p(0)}}(T_{p;k}) = \mathrm{jump}_{p(1)}(T_{p;k}) - 1$. In particular, when j is of Class $\mathcal{B}$ and $\varepsilon_j^\vee(T_{p;k}) = 2$, then $\varphi_j(p;k) = \mathrm{jump}_j(T_{p;k}) - 1 = \mathrm{jump}_j(T_{p;k}) - \delta_{j,p(0)}(1-\delta_{j,\mathrm{ext}_p^-})$. When j is of Class $\mathcal{D}$, (j,j') is a Class $\mathcal{D}$ pair, and $p(0) = j', p(1) = j$ then $\varphi_j^-(p;k) = -1$. Note, for these modules Remark 5.3 does not apply. Further, such p will not ever arise in our main theorems. See Remark 6.6 below as well as Remark 7.8.

Computing the values in Table 2 is an easy exercise using equation (118). Calculating jump_j is a key ingredient to calculating φ_j^Λ and from this one obtains Table 3. Then it is easy to check Remark 6.5 holds for $k > 1$.

Remark 6.6. Recall that a cyclotomic path p of tail weight $(\Lambda_{i_1}, \Lambda_{i_2})$ has a unique extension to its tail by an i_1-arrow, i.e. $\mathrm{ext}_p^- = \{i_1\}$. By Remark 6.5, this implies $\varphi^-(p;k) = \Lambda_{i_1}$. The condition on cyclotomic paths requires that $p(0) \neq p(1)$. When $k > 1$ this implies $\varphi_j(p;k) = \mathrm{jump}_j(T_{p;k})$ for all $j \in I$ and we shall use these two expressions interchangeably.

We now apply the lemmas and propositions from Section 5 to $T_{p;k}$.

Proposition 6.7. *Let p be a cyclotomic path of tail weight $(\Lambda_{p(-1)}, \Lambda_{p(0)})$ so that $\varphi^-(p;k) = \Lambda_{p(-1)}$. When $k \geq 1$ then the following hold:*

1. $T_{p;k} \in \mathrm{Rep}^{\Lambda_{p(0)}}$.
2. *If $j \neq p(-1)$ and $\varphi_j^+(p;k) = 0$, then*
$$\mathrm{pr}_{\Lambda_{p(0)}} \mathrm{Ind}\, T_{p;k} \boxtimes L(j) = \mathbf{0}. \tag{199}$$
3. *If $j \neq p(-1)$ but $\varphi_j^+(p;k) \geq 1$ then*
$$\mathrm{pr}_{\Lambda_{p(0)}} \mathrm{Ind}\, T_{p;k} \boxtimes L(j) \cong T_{p\star\pi(j);k+1}. \tag{200}$$

4. When $j = p(-1)$ and $\varphi_j^+(p;k) \geq 1$ then there is a surjection

$$\mathrm{pr}_{\Lambda_{p(0)}} \operatorname{Ind} T_{p;k} \boxtimes L(j) \twoheadrightarrow T_{p\star\pi(j);k+1} \tag{201}$$

   and for all composition factors K of $\operatorname{Ind} T_{p;k} \boxtimes L(j)$ such that $K \ncong T_{p\star\pi(j);k+1}$,

$$\varphi_j^{\Lambda_{p(0)}}(K) \leq \varphi_j(p;k) - 2. \tag{202}$$

Proof.

1. This follows from the definition of cyclotomic path and can also be verified by inspecting $\mathrm{Char}(T_{p;k})$ from Section 6.

2. This follows from Proposition 5.6.c.

3. If $p(-1) \neq j$ but $\varphi_j^+(p;k) \geq 1$, then either $\varphi_j(p;k) = 1$ or 2. When $\varphi_j(p;k) = 1$, the result is implied by Proposition 5.6.e, Remark 6.5 and Proposition 6.4. When $\varphi_j(p;k) = 2$ then j must be class $\mathcal{B}$ and the head of p can be extended twice by j-arrows. An examination of paths in $B^{1,1}$ whose head can be extended twice by class $\mathcal{B}$ arrows shows that $\varepsilon(T_{p;k}) = \Lambda_{p(k-1)}$, $a_{p(k-1),j} < 0$, and $\varphi_j(p;k-1) = 0$. Hence $\mathrm{pr}_{\Lambda_{p(0)}} \operatorname{Ind} T_{p;k-1} \boxtimes L(j) = \mathbf{0}$. The result then follows from Proposition 5.6.g and Proposition 6.4.

4. By Proposition 6.4 there is a surjection

$$\operatorname{Ind} T_{p;k} \boxtimes L(j) \twoheadrightarrow T_{p\star\pi(j);k+1} \tag{203}$$

   since $T_{p\star\pi(j);k+1} \cong \widetilde{f}_j T_{p;k}$. Because $\mathrm{pr}_{\Lambda_{p(0)}} T_{p\star\pi(j);k+1} = T_{p\star\pi(j);k+1}$, applying $\mathrm{pr}_{\Lambda_{p(0)}}$ to (203) gives us

$$\mathrm{pr}_{\Lambda_{p(0)}} \operatorname{Ind} T_{p;k} \boxtimes L(j) \twoheadrightarrow T_{p\star\pi(j);k+1}. \tag{204}$$

   By Proposition 4.6.2d if K is a composition factor of $\operatorname{Ind} T_{p;k} \boxtimes L(j)$ not isomorphic to $T_{p\star\pi(j);k+1}$ then $\varepsilon_j(K) \leq \varepsilon_j(T_{p;k})$ but $\mathrm{wt}_j(K) = \mathrm{wt}_j(T_{p;k}) - 2$. By formula (140) it follows that

$$\varphi_j^{\Lambda_{p(0)}}(K) \leq \varphi_j(p;k) - 2. \tag{205}$$

   It is possible $\mathrm{pr}_{\Lambda_{p(0)}} K = \mathbf{0}$; but (205) still holds even in this case.

□

7 Main theorems

We introduce some shortcut notation. For a simple $R(\nu)$-module A, define

$$\widetilde{e}^{\vee}_{i_k i_{k-1} \ldots i_2 i_1} A := \widetilde{e}^{\vee}_{i_k} \widetilde{e}^{\vee}_{i_{k-1}} \ldots \widetilde{e}^{\vee}_{i_2} \widetilde{e}^{\vee}_{i_1} A \tag{206}$$

and analogously,

$$\widetilde{f}^{\vee}_{i_k i_{k-1} \ldots i_2 i_1} A := \widetilde{f}^{\vee}_{i_k} \widetilde{f}^{\vee}_{i_{k-1}} \ldots \widetilde{f}^{\vee}_{i_2} \widetilde{f}^{\vee}_{i_1} A. \tag{207}$$

If p is a path in $B^{1,1}$ such that $p(0) = i_1$, $p(1) = i_2$, …, $p(k-1) = i_k$, we will also write

$$\widetilde{e}^{\vee}_{p;k} := \widetilde{e}^{\vee}_{i_k i_{k-1} \dots i_2 i_1} \qquad \text{and} \qquad \widetilde{f}^{\vee}_{p;k} := \widetilde{f}^{\vee}_{i_k i_{k-1} \dots i_2 i_1}. \tag{208}$$

Similarly we will occasionally write

$$T_{i_1 i_2 \dots i_k} := T_{p;k}, \tag{209}$$

particularly for small k.

7.1 Existence theorem

This is the first of our main theorems, which we term the "existence" theorem, and also sometimes refer to as a construction or algorithm.

Theorem 7.1. Let $A \in \mathrm{Rep}^{\Lambda_i}$ be a simple $R(\nu)$-module with $|\nu| \geq 1$ and i not a forbidden element of I.

1. There exists a cyclotomic path $p : \{0, 1, \dots, k-1\} \to I$ with $p(0) = i$ of tail weight $(\Lambda_{p(-1)}, \Lambda_i)$ and length k such that $\widetilde{e}^{\vee}_{p;k} A$ is a simple $R(\nu - \gamma_{p;k})$-module in $\mathrm{Rep}^{\Lambda_{p(-1)}}$.
2. Let $r(A) = k$ be the minimal k such that statement (1) holds and let $\mathcal{R}(A) = \widetilde{e}^{\vee}_{p;k} A$. Then there exists a surjection

$$\mathrm{pr}_{\Lambda_i} \operatorname{Ind} T_{p;k} \boxtimes \mathcal{R}(A) \twoheadrightarrow A. \tag{210}$$

Note by Remark 5.1 this is equivalent to proving $\operatorname{Ind} T_{p;k} \boxtimes \mathcal{R}(A) \twoheadrightarrow A$. We further conjecture the following.

Conjecture 7.2. Under the same hypotheses as Theorem 7.1

$$A \cong \operatorname{cosoc} \mathrm{pr}_{\Lambda_i} \operatorname{Ind} T_{p;k} \boxtimes \mathcal{R}(A). \tag{211}$$

Note that by Section 5.0.1, the crystal-theoretic consequence of Theorem 7.1 is a map $B(\Lambda_i) \to \bigoplus_j B^{1,1} \otimes B(\Lambda_j)$ given by

$$[A] \mapsto \enclose{circle}{p(k-1)} \otimes [\mathcal{R}(A)],$$

and where j runs over all possibilities for $p(-1)$. By abuse of notation, we let $\enclose{circle}{p(k-1)}$ be the node in $B^{1,1}$ that the path p ends at. In this way, each node of $B^{1,1}$ corresponds to the infinite collection of paths p that end at that node, and in turn to the collection of modules $T_{p;k}$. (As remarked in the introduction, this is not a categorification, but it is a useful correspondence.) If we choose to specify $B(\Lambda_i)$ as above, that further specifies that of that collection, we care about the p with $p(0) = i$.

To recover the crystal isomorphism (21) or (22) from Theorem 7.1 one must actually fix $p(-1)$ and let i vary (whereas the theorem fixes i). In many types X_ℓ, $i \in I$, specifying $p(0) = i$ determines $p(-1)$ (in particular when Λ_i is of level 1 and $B^{1,1}$ is perfect). In type A the relationship between (21) and (210) is transparent. [33] discusses the crystal isomorphism in type A in more depth. Examples 2.2, 7.16 and Remark 7.9 discuss a case when there are two choices for $p(-1)$, even when p is a cyclotomic path. Note that $p(0)$ and $p(-1)$ must be "adjacent" arrows in $B^{1,1}$. At the moment, the above theorem just gives us a map of nodes. Our second main theorem in Section 7.2 below will show that it is a morphism of crystals.

We will construct the path p in the proof of Theorem 7.1. We shall see p is not always unique. For our choice of path p we will also construct intermediate modules $\mathcal{R}_t(A) = \widetilde{e}^{\vee}_{p;t}A$ and refer to this process as our *algorithm.* We first prove a series of lemmas.

Lemma 7.3. Suppose that $A \in \mathrm{Rep}^{\Lambda_i}$ is a simple $R(\nu)$-module, D is a simple $R(\nu - \gamma_{p;k})$-module, p is a cyclotomic path of length k and tail weight $(\Lambda_{p(-1)}, \Lambda_i)$, and there is a surjection

$$\operatorname{Ind} T_{p;k} \boxtimes D \twoheadrightarrow A. \tag{212}$$

Then $D \in \mathrm{Rep}^{\Lambda_{p(-1)}+\varphi^+(p;k)}$.

Proof. By Remark 6.6, $\varphi^-(p;k) = \Lambda_{p(-1)}$ so in particular $\Lambda_{p(-1)} + \varphi^+(p;k) = \varphi(p;k)$. Proposition 5.6.d then implies the result. ☐

Lemma 7.4. Let p be a cyclotomic path of length k and tail weight $(\Lambda_{p(-1)}, \Lambda_i)$. Suppose that $A \in \mathrm{Rep}^{\Lambda_i}$ is a simple $R(\nu)$-module, D is a simple $R(\nu - \gamma_{p;k})$-module and there is a surjection

$$\operatorname{Ind} T_{p;k} \boxtimes D \twoheadrightarrow A. \tag{213}$$

If $j \in \mathrm{ext}_p^+$ such that $\varepsilon_j^{\vee}(D) \neq 0$ and either $j \neq p(-1)$ or $\varepsilon_j^{\vee}(D) = \varphi_j(p;k)$, then there is a surjection

$$\operatorname{Ind} T_{p\star\pi(j);k+1} \boxtimes \widetilde{e}_j^{\vee} D \twoheadrightarrow A. \tag{214}$$

Proof. Since $\varepsilon_j^{\vee}(D) \neq 0$, there is a surjection

$$\operatorname{Ind} L(j) \boxtimes \widetilde{e}_j^{\vee} D \twoheadrightarrow D. \tag{215}$$

Then by the exactness of induction we get

$$\operatorname{Ind} T_{p;k} \boxtimes L(j) \boxtimes \widetilde{e}_j^{\vee} D \twoheadrightarrow A. \tag{216}$$

First suppose that $p(-1) \neq j$. Then by Proposition 6.7.3

$$\mathrm{pr}_{\Lambda_i} \operatorname{Ind} T_{p;k} \boxtimes L(j) \cong T_{p\star\pi(j);k+1} \tag{217}$$

and by Proposition 5.6.f applying pr_{Λ_i} to (216) gives

$$\operatorname{Ind} T_{p\star\pi(j);k+1} \boxtimes \widetilde{e}_j^{\vee} D \twoheadrightarrow \mathrm{pr}_{\Lambda_i} \operatorname{Ind} T_{p\star\pi(j);k+1} \boxtimes \widetilde{e}_j^{\vee} D \twoheadrightarrow \mathrm{pr}_{\Lambda_i} A \cong A. \tag{218}$$

Second, if $j = p(-1)$ then both $\varphi^-(p;k)$ and $\varphi^+(p;k)$ have Λ_j in their support so that $\varepsilon_j^{\vee}(D) = \varphi_j(p;k) \geq 2$. By Proposition 6.4, $\mathrm{pr}_{\Lambda_i} \operatorname{Ind} T_{p;k} \boxtimes L(j)$ has cosocle isomorphic to $T_{p\star\pi(j);k+1}$. So if (218) doesn't hold then by Remark 4.5 there exists some other composition factor K of $\operatorname{Ind} T_{p;k} \boxtimes L(j)$ such that

$$\operatorname{Ind} K \boxtimes \widetilde{e}_j^{\vee} D \twoheadrightarrow A. \tag{219}$$

By Proposition 6.7.4, K satisfies $\varphi_j^{\Lambda_i}(K) = \varphi_j^{\Lambda_i}(T_{p;k}) - 2 =: d$. Observe that

$$\varepsilon_j^\vee(\widetilde{e}_j^\vee D) = \varepsilon_j^\vee(D) - 1 = \varphi_j(p;k) - 1 = d + 1 > \varphi_j^{\Lambda_i}(K), \tag{220}$$

which implies that

$$\mathrm{pr}_{\Lambda_i} \operatorname{Ind} K \boxtimes L(j^{d+1}) \boxtimes (\widetilde{e}_j^\vee)^{d+2} D \twoheadrightarrow A. \tag{221}$$

But $\mathrm{pr}_{\Lambda_i} \operatorname{Ind} K \boxtimes L(j^{d+1}) \boxtimes (\widetilde{e}_j^\vee)^{d+2} D = \mathbf{0}$ by Proposition 5.6.c. Hence the surjection (216) must factor through $\operatorname{Ind} T_{p\star\pi(j);k+1} \boxtimes \widetilde{e}_j^\vee D$, i.e. (214) holds. ☐

Lemma 7.5. Suppose i is of class $\mathcal{B}$ and that locally in $B^{1,1}$



Set $\Lambda = \Lambda_i + \Lambda_h$ and suppose that $D \in \mathrm{Rep}^\Lambda$ is simple, $\widetilde{e}_h^\vee D \in \mathrm{Rep}^{2\Lambda_i}$, and $\widetilde{e}_h^\vee D \notin \mathrm{Rep}^{\Lambda_i}$. Then

1. $\varepsilon^\vee(\widetilde{e}_{ih}^\vee D) = \Lambda_i$,
2. There exists a surjection
$$\operatorname{Ind} T_{hi} \boxtimes \widetilde{e}_{ih}^\vee D \twoheadrightarrow D \tag{222}$$

Proof. 1. As $\varepsilon^\vee(\widetilde{e}_h^\vee D) = 2\Lambda_i$ there is a surjection

$$\operatorname{Ind} L(i) \boxtimes \widetilde{e}_{ih}^\vee D \twoheadrightarrow \widetilde{e}_h^\vee D. \tag{223}$$

Because i is a class $\mathcal{B}$ node, $a_{ij} = 0$ unless $j = h$ or $j = i$ in which case $a_{hi} = -1$ or $a_{ii} = 2$ respectively. Using equation (140), $\varphi^{2\Lambda_i}(L(i)) = \Lambda_i + \Lambda_h$, and it follows from Proposition 5.6.c that $\widetilde{e}_{ih}^\vee D \in \mathrm{Rep}^\Lambda$. We therefore only need to show that $\varepsilon_h^\vee(\widetilde{e}_{ih}^\vee D) = 0$.

There exist $m_i, m_h \in \mathbb{Z}_{>0}$, a simple $R(m_i\alpha_i + m_h\alpha_h)$-module $K \in \mathrm{Rep}^\Lambda$, and a simple $R(\nu - (m_i\alpha_i + m_h\alpha_h))$-module N such that $\varepsilon_i^\vee N = \varepsilon_h^\vee N = 0$, and a surjection

$$\operatorname{Ind} K \boxtimes N \twoheadrightarrow D. \tag{224}$$

Then $K \in \mathrm{Rep}^\Lambda$, $\widetilde{e}_h^\vee K \in \mathrm{Rep}^{2\Lambda_i}$, and $\widetilde{e}_h^\vee K \notin \mathrm{Rep}^{\Lambda_i}$. An analysis of the classification of simple modules in $\mathrm{Rep}^{\Lambda_i+\Lambda_h}$ in the Appendix shows that K must be isomorphic to one of the following:

(a) $\mathcal{L}(hii) \cong \widetilde{f}_h^\vee \widetilde{f}_i^\vee L(i)$,

(b) $\widetilde{f}_h \mathcal{L}(hii) \cong \widetilde{f}_h^\vee \widetilde{f}_i^\vee \mathcal{L}(ih)$, which has character $(1 + q_i^2)[hiih]$ (up to overall grading shift),

(c) $\operatorname{Ind} \mathcal{L}(hi) \boxtimes \mathcal{L}(ihi) \cong \widetilde{f}_h^\vee \widetilde{f}_i^\vee \mathcal{L}(ihi)$.

By Proposition 4.10, it follows that there is a surjection,

$$\operatorname{Ind} \widetilde{e}_{ih}^\vee K \boxtimes N \twoheadrightarrow \widetilde{e}_{ih}^\vee D. \tag{225}$$

From the three cases above, $\widetilde{e}_{ih}^\vee K$ is isomorphic to either $L(i)$, $\mathcal{L}(ih)$, or $\mathcal{L}(ihi)$. In all three cases $\varepsilon_h^\vee(\widetilde{e}_{ih}^\vee K) = 0$. Since $\varepsilon_h^\vee(N) = 0$, it follows from the Shuffle Lemma that $\varepsilon_h^\vee(\widetilde{e}_{hi}^\vee D) = 0$.

2. There is a nonzero map
$$\operatorname{Ind} L(h) \boxtimes L(i) \boxtimes \widetilde{e}^{\vee}_{ih} D \twoheadrightarrow D. \tag{226}$$
This surjection factors through either $\operatorname{Ind} \mathcal{L}(ih) \boxtimes \widetilde{e}^{\vee}_{ih} D$ or $\operatorname{Ind} \mathcal{L}(hi) \boxtimes \widetilde{e}^{\vee}_{ih} D$. Since $\varepsilon^{\vee}_{h}(D) = 1$ but $\varepsilon^{\vee}_{h}(\mathcal{L}(ih)) = 0$ and $\varepsilon^{\vee}_{h}(\widetilde{e}^{\vee}_{ih} D) = 0$ by part (1), then $e^{\vee}_{h}(D) \neq \mathbf{0}$ but $e^{\vee}_{h}(\operatorname{Ind} \mathcal{L}(ih) \boxtimes \widetilde{e}^{\vee}_{hi} D) = \mathbf{0}$ a contradiction to the exactness of $e^{\vee}_{h}$. It follows that (226) factors through
$$\operatorname{Ind} \mathcal{L}(hi) \boxtimes \widetilde{e}^{\vee}_{ih} D \twoheadrightarrow D. \tag{227}$$

□

Lemma 7.6. Let i belong to class $\mathcal{B}$ such that locally in $B^{1,1}$



Suppose that p is a cyclotomic path in $B^{1,1}$ of tail weight (Λ_i, Λ_i) and length $k+2$ such that $p(k-2) = h'$, $p(k-1) = h$, $p(k) = p(k+1) = i$. Suppose further that A is a simple $R(\nu)$-module in $\operatorname{Rep}^{\Lambda_i}$, D is a simple $R(\nu - \gamma_{p;k})$-module such that $D \in \operatorname{Rep}^{2\Lambda_i}$, $D \notin \operatorname{Rep}^{\Lambda_i}$, and $\widetilde{f}^{\vee}_{h} D \in \operatorname{Rep}^{\Lambda_i + \Lambda_h}$, and there are surjections
$$\operatorname{Ind} T_{p;k} \boxtimes D \twoheadrightarrow A, \tag{228}$$
$$\operatorname{Ind} T_{p;k-1} \boxtimes \widetilde{f}^{\vee}_{h} D \twoheadrightarrow A. \tag{229}$$
Then $\widetilde{e}^{\vee}_{i} D \in \operatorname{Rep}^{\Lambda_i}$ and there is a surjection
$$\operatorname{Ind} T_{p;k+1} \boxtimes \widetilde{e}^{\vee}_{i} D \twoheadrightarrow A \tag{230}$$

Proof. The claim that $\widetilde{e}^{\vee}_{i} D \in \operatorname{Rep}^{\Lambda_i}$ follows immediately from Lemma 7.5 and the fact that $\widetilde{f}^{\vee}_{h} D \in \operatorname{Rep}^{\Lambda_i + \Lambda_h}$, $D \in \operatorname{Rep}^{2\Lambda_i}$, $D \notin \operatorname{Rep}^{\Lambda_i}$.

To construct the surjection (230) we first prove that
$$\operatorname{pr}_{\Lambda_i} \operatorname{Ind} T_{p;k-1} \boxtimes T_{hi} \cong T_{p;k+1}. \tag{231}$$

Recall the notation $T_{hi} = \mathcal{L}(hi)$. Examining Section 6, we see $\varepsilon_h(T_{p;k+1}) = 0$, so recalling Proposition 4.9 the map
$$\operatorname{Ind} T_{p;k-1} \boxtimes L(h) \boxtimes L(i) \twoheadrightarrow T_{p;k+1}$$
cannot factor through $\operatorname{Ind} T_{p;k-1} \boxtimes \mathcal{L}(ih)$ yielding surjections
$$\operatorname{Ind} T_{p;k-1} \boxtimes L(h) \boxtimes L(i) \twoheadrightarrow \operatorname{Ind} T_{p;k-1} \boxtimes T_{hi} \twoheadrightarrow T_{p;k+1}. \tag{232}$$

Since $i = p(0) = p(-1) \neq h$, Proposition 5.6.f, Proposition 6.7.3, the fact $\operatorname{pr}_{\Lambda_i} T_{p;k+1} \cong T_{p;k+1}$, and the right exactness of $\operatorname{pr}_{\Lambda}$ imply we have
$$\begin{tikzcd} \operatorname{pr}_{\Lambda_i} \operatorname{Ind} T_{p;k-1} \boxtimes L(h) \boxtimes L(i) \arrow[r, two heads] \arrow[d, phantom, sloped, "\cong"] & \operatorname{pr}_{\Lambda_i} \operatorname{Ind} T_{p;k-1} \boxtimes T_{hi} \arrow[r, two heads] & T_{p;k+1}. \\ \operatorname{pr}_{\Lambda_i} \operatorname{Ind} T_{p;k} \boxtimes L(i) & & \end{tikzcd} \tag{233}$$

Keeping in mind the lower bounds on ℓ and by inspection of the types X_ℓ that have class $\mathcal{B}$ nodes, we see that (h,h') do *not* form a class $\mathcal{D}$ pair. Hence all composition factors $K \not\cong \operatorname{cosoc}(\operatorname{Ind} T_{p;k} \boxtimes L(i)) \cong T_{p;k+1}$ of $\operatorname{pr}_{\Lambda_i} \operatorname{Ind} T_{p;k} \boxtimes L(i)$ have $\varepsilon(K) = \Lambda_h$. As $e_h(\operatorname{Ind} T_{p;k-1} \boxtimes T_{hi}) = \mathbf{0}$ it follows that

$$\operatorname{pr}_{\Lambda_i} \operatorname{Ind} T_{p;k-1} \boxtimes T_{hi} \cong T_{p;k+1}. \tag{234}$$

By Lemma 7.5 there is a surjection

$$\operatorname{Ind} T_{hi} \boxtimes \widetilde{e}_i^{\vee} D \twoheadrightarrow \widetilde{f}_h^{\vee} D. \tag{235}$$

The exactness of induction and (229) then give

$$\operatorname{Ind} T_{p;k-1} \boxtimes T_{hi} \boxtimes \widetilde{e}_i^{\vee} D \twoheadrightarrow A. \tag{236}$$

By (234), Proposition 5.6.f implies that

$$\operatorname{pr}_{\Lambda_i} \operatorname{Ind} T_{p;k+1} \boxtimes \widetilde{e}_i^{\vee} D \twoheadrightarrow A. \tag{237}$$

□

We now prove our first main theorem, Theorem 7.1.

Proof. Since i is not a forbidden element of I then Lemma 3.3 states that we can find at least one cyclotomic path of length 1 and tail weight $(\Lambda_{p(-1)}, \Lambda_i)$ in $B^{1,1}$ for some $p(-1) \in I$. Choose one of these cyclotomic paths and call it p_1. We will construct the path p in the statement of the theorem by repeatedly extending the head of p_1. In particular p_t will be the length t path that results from extending the head of p_{t-1} by some $j \in \operatorname{ext}^+_{p_{t-1}}$ so that $p_t := p_{t-1} \star \pi(j)$. Since the tail is fixed for $p_1, p_2, \ldots, p_t, \ldots$, all p_t will be cyclotomic paths of tail weight $(\Lambda_{p(-1)}, \Lambda_i)$, and therefore for all $t \geq 1$, $\varphi^-(p_t; t) = \Lambda_{p(-1)}$. When i is part of a class $\mathcal{D}$ pair (i,j) then $p(-1) = j$, and we add the requirement to p_t that we *favor* extension by i-arrows over extension by j-arrows. In other words when $\operatorname{ext}^+_{p_t} = \{i, j\}$ set

$$p_{t+1} = p_t \star \pi(i) \quad \text{and} \quad p_{t+2} = p_{t+1} \star \pi(j) \tag{238}$$

unless the algorithm terminates before $t+2$.

Set $\mathcal{R}_0(A) := A$, and

$$\mathcal{R}_t(A) := \widetilde{e}^{\vee}_{p_t(t-1)p_t(t-2)\ldots p_t(1)p_t(0)} A = \widetilde{e}^{\vee}_{p_t;t} A. \tag{239}$$

We will show inductively that there exist surjections

$$\operatorname{pr}_{\Lambda_i} \operatorname{Ind} T_{p_t;t} \boxtimes \mathcal{R}_t(A) \twoheadrightarrow A. \tag{240}$$

For each of these surjections Lemma 7.3 implies that $\mathcal{R}_t(A) \in \operatorname{Rep}^{\Lambda_{p(-1)} + \varphi^+(p_t;t)}$. The induction will end at the smallest k such that $\mathcal{R}_k(A) \in \operatorname{Rep}^{\Lambda_{p(-1)}}$. Then we set $p = p_k$, $r(A) = k$, and $\mathcal{R}(A) = \mathcal{R}_k(A)$.

In the base case $t = 0$, $\mathcal{R}_0(A) = A$. If $|\nu| = 1$, then $A = L(i)$ and $\widetilde{e}^{\vee}_{p_1(0)} L(i) \cong \mathbb{1} \in \operatorname{Rep}^{\Lambda_{p(-1)}}$, so $r(A) = 1$. The existence of the surjection in this case is vacuous.

Assume that $|\nu| > 1$. Then $\mathcal{R}_1(A) = \widetilde{e}^{\vee}_{p_1(0)}A \neq \mathbf{0}, \mathbb{1}$. By Proposition 4.6.2 there is a surjection

$$\operatorname{Ind} T_{p_1(0)} \boxtimes \mathcal{R}_1(A) \cong \operatorname{Ind} L(p_1(0)) \boxtimes \widetilde{e}^{\vee}_{p_1(0)}A \twoheadrightarrow A. \tag{241}$$

As noted above $\mathcal{R}_1(A) \in \operatorname{Rep}^{\varphi^+(p_1;1)+\Lambda_{p(-1)}}$.

Now suppose that we have shown that for $1 \leq t \leq r$ there exist surjections

$$\operatorname{pr}_{\Lambda_i} \operatorname{Ind} T_{p_t;t} \boxtimes \mathcal{R}_t(A) \twoheadrightarrow A \tag{242}$$

with $\mathcal{R}_t(A) \in \operatorname{Rep}^{\Lambda_{p(-1)}+\varphi^+(p_t;t)}$ but for $t < r$, $\mathcal{R}_t(A) \notin \operatorname{Rep}^{\Lambda_{p(-1)}}$. If in fact $\mathcal{R}_r(A) \in \operatorname{Rep}^{\Lambda_{p(-1)}}$ then we are done and we set $p_r = p$, $r(A) = r$, and $\mathcal{R}(A) = \mathcal{R}_r(A)$. If not, then there is some $j \in I$ with Λ_j in the support of $\varphi^+(p_r; r)$ such that either $j \neq p(-1)$ and $\varepsilon^{\vee}_j(\mathcal{R}_r(A)) \geq 1$, or $j = p(-1)$ and $\varepsilon^{\vee}_{p(-1)}(\mathcal{R}_r(A)) \geq 2$. By Remark 6.5 in either case we can extend the head of p_r by a j-arrow and set $p_{r+1} := p_r \star \pi(j)$. p_{r+1} is a cyclotomic path of tail weight $(\Lambda_{p(-1)}, \Lambda_i)$ and length $r+1$. Recall that if $(p(-1), i)$ is a class $\mathcal{D}$ pair, then we always favor extension by i-arrows. Set $\mathcal{R}_{r+1}(A) := \widetilde{e}^{\vee}_{p_{r+1}(r)}\mathcal{R}_r(A) = \widetilde{e}^{\vee}_j \mathcal{R}_r(A)$. When $j \neq p(-1)$, Lemma 7.4 implies that there is a surjection

$$\operatorname{pr}_{\Lambda_i} \operatorname{Ind} T_{p_{r+1};r+1} \boxtimes \mathcal{R}_{r+1}(A) \twoheadrightarrow A, \tag{243}$$

and then Lemma 7.3 and (243) imply that $\mathcal{R}_{r+1}(A) \in \operatorname{Rep}^{\Lambda_{p(-1)}+\varphi^+(p_{r+1};r+1)}$.

If $j = p(-1)$ and $\varepsilon^{\vee}_{p(-1)}(\mathcal{R}_r(A)) > 1$, then from the formulas for $\varphi^+(p; r)$ in Table 4, $\varepsilon^{\vee}_{p(-1)}(\mathcal{R}_r(A)) \leq \varphi^+_{p(-1)}(p; r) + 1 \leq 3$. If $\varepsilon^{\vee}_{p(-1)}(\mathcal{R}_r(A)) = 3$ then $\varepsilon^{\vee}_{p(-1)}(\mathcal{R}_r(A)) = \varphi_{p(-1)}(p_r; r)$. In this case Lemma 7.4 implies that surjection (243) exists and as before $\mathcal{R}_{r+1}(A) \in \operatorname{Rep}^{\Lambda_{p(-1)}+\varphi^+(p_{r+1};r+1)}$.

The only remaining case is when $\varepsilon^{\vee}_{p(-1)}(\mathcal{R}_r(A)) = 2$, and either $\varphi_{p(-1)}(p_r; r) = 2$ or $\varphi_{p(-1)}(p_r; r) = 3$. When the former holds again Lemma 7.4 gives surjection (243) and hence $\mathcal{R}_{r+1}(A) \in \operatorname{Rep}^{\Lambda_{p(-1)}+\varphi^+(p_{r+1};r+1)}$. If $\varphi_{p(-1)}(p_r; r) = 3$, then because p_{r+1} is a cyclotomic path, $j = p(-1) = p(0) = i$ is a class $\mathcal{B}$ node, $p_r(r-1) \neq p(-1)$ and $r \geq 4$. Because $r \geq 4$ the inductive hypothesis implies that there is a surjection

$$\operatorname{Ind} T_{p_r;r-1} \boxtimes \mathcal{R}_{r-1}(A) \twoheadrightarrow A \tag{244}$$

(here we make the identification $T_{p_{r-1};r-1} \cong T_{p_r;r-1}$ as $p_r(t) = p_{r-1}(t)$ for all $0 \leq t < r-1$). Then Lemma 7.6 implies that there is a surjection

$$\operatorname{pr}_{\Lambda_i} \operatorname{Ind} T_{p_{r+1};r+1} \boxtimes \mathcal{R}_{r+1}(A) \twoheadrightarrow A \tag{245}$$

and $\mathcal{R}_{r+1}(A) \in \operatorname{Rep}^{\Lambda_{p(-1)}}$ so $\mathcal{R}(A) := \mathcal{R}_{r+1}(A)$ and $r(A) = r+1$ and the induction terminates. This proves the inductive step.

We continue the induction until we reach the first k such that $\mathcal{R}_k(A) \in \operatorname{Rep}^{\Lambda_{p(-1)}}$. At this point we set $p = p_k$, $r(A) = k$, $\mathcal{R}(A) = \mathcal{R}_k(A)$. $\mathcal{R}_k(A)$ is a $R(\nu - \gamma_{p_k;k})$-module. As $|\nu - \gamma_{p;k}| = |\nu| - k$, the algorithm necessarily terminates at or before $k = |\nu|$. If $k = |\nu|$ then $\mathcal{R}_{|\nu|}(A) = \mathbb{1} \in \operatorname{Rep}^{\Lambda_{p(-1)}}$. In this case

$$T_{p_{|\nu|};|\nu|} \cong \operatorname{pr}_{\Lambda_i} \operatorname{Ind} T_{p_{|\nu|};|\nu|} \boxtimes \mathbb{1} \twoheadrightarrow A \tag{246}$$

so that in fact $A \cong T_{p_{|\nu|};|\nu|}$.

Note that for all $r \leq r(A)$ and $t < r$, $p(t) = p_r(t)$ so all intermediate paths constructed above are compatible.

$\square$

In Theorem 7.1 we chose $p = p_k$ such that whenever (i, j) is a class $\mathcal{D}$ pair and $\{p(t-1), p(t)\} = \{i, j\}$ we had $p(t-1) = i$, $p(t) = j$. However if p' agree with p except that for $t \in \eta^i_j(\underline{\tau_{\mathbf{p;k}}})$ we allow $\{p'(t-1), p'(t)\} = \{i, j\}$ in either order then in fact $T_{p;k} \cong T_{p';k}$ even though $p \neq p'$. This also applies to class $\mathcal{D}$ pairs (j, j') when $i \notin \{j, j'\}$. Here p and p' take different routes around the bifurcation in $B^{1,1}$ generated by the class $\mathcal{D}$ pair (i, j). (Of course if $p(k-2) \notin \{i, j\}$ and $p(k-1) = i$ we must also have $p'(k-1) = i$ since p ends before it traverses the entire bifurcation). This phenomenon only occurs in types $D^{(1)}_\ell$, $B^{(1)}_\ell$, and $A^{(2)}_{2\ell-1}$. (See Sections 2.3 and 6.)

The following proposition shows that choosing route p' over p does not affect the output of the algorithm from Theorem 7.1. In other words, after the initial choice of length 1 cyclotomic path p_1, $\mathcal{R}(A)$ and $r(A)$ are well-defined. Part 1 of Proposition 7.7 shows that $\widetilde{e}^\vee_{p;k} A = \widetilde{e}^\vee_{p';k} A$ while part 2 in conjunction with part 1 shows that $r(A)$ is well-defined. It follows from this that in fact $\mathcal{R}(A)$ is well-defined.

Proposition 7.7. Suppose that locally in $B^{1,1}$



so that (j, j') is a class $\mathcal{D}$ pair. Let $A \in \mathrm{Rep}^{\Lambda_i}$ be a simple $R(\nu)$-module, and suppose r steps of the algorithm from Theorem 7.1 have been executed so we have constructed a simple $R(\nu - \gamma_{p;r})$-module $\mathcal{R}_r(A)$, a cyclotomic path p_r with length r and tail weight $(\Lambda_{p(-1)}, \Lambda_i)$, and a surjection

$$\operatorname{Ind} T_{p_r;r} \boxtimes \mathcal{R}_r(A) \twoheadrightarrow A \tag{247}$$

Finally, suppose that $p_r(r-1) = h$, $\varphi^+(p_r; r) = \Lambda_j + \Lambda_{j'}$. Then

1. $\widetilde{e}^\vee_{j'} \widetilde{e}^\vee_j \mathcal{R}_r(A) \cong \widetilde{e}^\vee_j \widetilde{e}^\vee_{j'} \mathcal{R}_r(A)$.

2. If $\widetilde{e}^\vee_j \mathcal{R}_r(A) \in \mathrm{Rep}^{\Lambda_{p(-1)}}$ then $\mathcal{R}_r(A) \in \mathrm{Rep}^{\Lambda_{p(-1)} + \Lambda_j}$.

Proof. Part 1 follows directly from Proposition 4.19 and the fact that for class $\mathcal{D}$ pairs (j, j'), $a_{jj'} = 0$. Part 2 is a direct application of part 1 after noting that by Lemma 7.3, $\mathcal{R}_r(A) \in \mathrm{Rep}^{\Lambda_{p(-1)} + \Lambda_j + \Lambda_{j'}}$. $\square$

Remark 7.8. Let (j, j') be a class $\mathcal{D}$ pair. The attentive reader will notice that the case where $\mathcal{R}_r(A) \cong \operatorname{Ind} L(j) \boxtimes L(j')$ with $p(-1) = j$ and $i = j'$ could potentially give two different values of $r(A)$ depending on whether one chooses to extend p_r by j' or j. In the former case $\mathcal{R}(A) = \mathcal{R}_{r+1}(A) = L(j) \in \mathrm{Rep}^{\Lambda_{p(-1)}}$ and $r(A) = r + 1$. In the latter case $\mathcal{R}_{r+1}(A) = L(j') \notin \mathrm{Rep}^{\Lambda_{p(-1)}}$ and the algorithm tells us to continue to get $\mathcal{R}(A) = \mathcal{R}_{r+2}(A) = \mathbb{1}$, yielding

$r(A) = r+2$. Recall that when $(i, p(-1))$ is a class $\mathcal{D}$ pair, then the algorithm requires that when building our path p, we always favor extending by i-arrows so that actually, Theorem 7.1 will always give $\mathcal{R}(A) \cong L(j)$ and $r(A) = r+1$ in this case. We require this favoring precisely to avoid the above situation.

7.2 The action of the crystal operators

Next we study the action of the crystal operators $\widetilde{e}_j$ and $\widetilde{f}_j$ on (210) to show that the map in part 2 of Theorem 7.1 categorifies our crystal isomorphism $\mathcal{T}$.

Remark 7.9. In some types, having chosen $p(0) = i$, the choice of $p(1)$ and consequently p_t, $t < r(A)$ is forced upon us. In other types, such as $C_\ell^{(1)}$, there can be 2 choices for $p(1)$ (and hence $p(-1)$). This choice is mirrored by the combinatorial structure of $B^{1,1} \otimes B(\Lambda_i)$. See Example 2.2 and Example 7.16.

Compare the theorems below with the crystal-theoretic statements (17) and (18). As in [23] simple modules in Rep^{Λ_i} correspond to nodes in the highest weight crystal $B(\Lambda_i)$. Each node b of the KR crystal $B^{1,1}$ (respectively $B^{\ell,1}$ for type $A_\ell^{(1)}$) corresponds to an infinite family of $R(\gamma_{p;k})$-modules $T_{p;k}, k \in \mathbb{Z}_{\geq 0}$ that satisfy $\varepsilon(T_{p;k}) = \varepsilon(b)$. It is in this manner that the main theorems of this paper give a categorification of the crystal isomorphism $\mathcal{T}$.

Theorem 7.10. *Let $A \in \mathrm{Rep}^{\Lambda_i}$ be a simple $R(\nu)$-module and $j \in I$ be such that $\widetilde{e}_j A \neq \mathbf{0}$. When i is class $\mathcal{B}$ we furthermore require that $|\nu| > 1$. Let p be a cyclotomic path of tail weight $(\Lambda_{p(-1)}, \Lambda_i)$ and length $k = r(A)$, and $\mathcal{R}(A) = \widetilde{e}^\vee_{p;k} A$, as constructed by the algorithm in Theorem 7.1. Then there exists a surjection*

$$\operatorname{Ind} \widetilde{e}_j T_{p;k} \boxtimes \mathcal{R}(A) \twoheadrightarrow \widetilde{e}_j A \quad \text{if} \quad \varepsilon_j(T_{p;k}) > \varphi_j^{\Lambda_{p(-1)}}(\mathcal{R}(A)), \tag{248}$$

$$\operatorname{Ind} T_{p;k} \boxtimes \widetilde{e}_j \mathcal{R}(A) \twoheadrightarrow \widetilde{e}_j A \quad \text{if} \quad \varepsilon_j(T_{p;k}) \leq \varphi_j^{\Lambda_{p(-1)}}(\mathcal{R}(A)). \tag{249}$$

Theorem 7.11. *Let i, p, k, A, and $\mathcal{R}(A)$ be as in Theorem 7.1. Let $j \in I$ be such that $\mathrm{pr}_{\Lambda_i} \widetilde{f}_j A \neq \mathbf{0}$. Then there exists a surjection*

$$\operatorname{Ind} \widetilde{f}_j T_{p;k} \boxtimes \mathcal{R}(A) \twoheadrightarrow \widetilde{f}_j A \quad \text{if} \quad \varepsilon_j(T_{p;k}) \geq \varphi_j^{\Lambda_{p(-1)}}(\mathcal{R}(A)), \tag{250}$$

$$\operatorname{Ind} T_{p;k} \boxtimes \widetilde{f}_j \mathcal{R}(A) \twoheadrightarrow \widetilde{f}_j A \quad \text{if} \quad \varepsilon_j(T_{p;k}) < \varphi_j^{\Lambda_{p(-1)}}(\mathcal{R}(A)). \tag{251}$$

Theorem 7.11 follows directly from Theorem 7.10, therefore we will only prove 7.10.

Remark 7.12. In the case $\varepsilon_j(T_{p;k}) \geq \varphi_j^{\Lambda_{p(-1)}}(\mathcal{R}(A))$, $\mathrm{pr}_{\Lambda_i} \widetilde{f}_j A \neq 0$ it will hold that $\widetilde{f}_j T_{p;k} \cong T_{p\star\pi(j);k+1}$ and $r(\widetilde{f}_j A) = r(A)+1$. (This relies in part on the properties of cyclotomic paths.)

We state a few lemmas below that follow directly from propositions in Section 4.6.

Lemma 7.13. *Let M be a simple $R(\nu)$-module and $h \neq j$. If $\widetilde{e}_j M \neq \mathbf{0}$ then $\varepsilon_h^\vee(\widetilde{e}_j M) = \varepsilon_h^\vee(M)$.*

Proof. This follows from Proposition 4.17.1d, and the definition of $\varepsilon_h^\vee(M)$. $\square$

Lemma 7.14. Suppose that D is a simple $R(\nu)$-module. If $\widetilde{e}_jD \neq \mathbf{0}$ and $\mathrm{jump}_j(D) \geq 1$, then $\varepsilon_j^\vee(D) = \varepsilon_j^\vee(\widetilde{e}_jD)$.

Proof. Set $N = \widetilde{e}_jD$. By (117) $\mathrm{jump}_j(N) = \mathrm{jump}_j(\widetilde{f}_jN) + 1 \geq 2$. Proposition 4.14 then implies, $\varepsilon_j^\vee(N) = \varepsilon_j^\vee(\widetilde{f}_jN)$, which is equivalent to $\varepsilon_j^\vee(\widetilde{e}_jD) = \varepsilon_j^\vee(D)$. □

In the following lemma we allow $i = j$.

Lemma 7.15. Let $i, j \in I$. Suppose that $D \in \mathrm{Rep}^{\Lambda_i}$ is a simple $R(\nu)$-module and $\widetilde{e}_jD \neq \mathbf{0}$.

1. If $\mathrm{jump}_j(D) \geq 1$, then
   (a) $\varepsilon_h^\vee(\widetilde{e}_jD) = \varepsilon_h^\vee(D)$ for all $h \in I$,
   (b) $\widetilde{e}_jD \in \mathrm{Rep}^{\Lambda_i}$,
   (c) $\varepsilon_j^\vee(\widetilde{e}_j\widetilde{f}_j^\vee D) = \varepsilon_j^\vee(\widetilde{f}_j^\vee D)$
2. If $\mathrm{jump}_j(D) \geq 2$, then furthermore $\varepsilon_j^\vee(\widetilde{e}_j\widetilde{f}_j^\vee\widetilde{f}_j^\vee D) = \varepsilon_j^\vee(\widetilde{f}_j^\vee\widetilde{f}_j^\vee D)$.

Proof. 1. When $h \neq j$ the equality $\varepsilon_h^\vee(\widetilde{e}_jD) = \varepsilon_h^\vee(D)$ holds by Lemma 7.13, and when $h = j$ it holds by Lemma 7.14. Since $\varepsilon_h^\vee(\widetilde{e}_jD) = \varepsilon_h^\vee(D)$ for all $h \in I$ then by Proposition 5.2, $\widetilde{e}_jD \in \mathrm{Rep}^{\Lambda_i}$.

Since $\mathrm{jump}_j(\widetilde{e}_jD) \geq 2$ then by Proposition 4.17.2c $\widetilde{e}_j\widetilde{f}_j^\vee D \cong \widetilde{f}_j^\vee\widetilde{e}_jD$. With the equality $\varepsilon_j^\vee(\widetilde{e}_jD) = \varepsilon_j^\vee(D)$ this implies that

$$\varepsilon_j^\vee(\widetilde{e}_j\widetilde{f}_j^\vee D) = \varepsilon_j^\vee(\widetilde{f}_j^\vee\widetilde{e}_jD) = \varepsilon_j^\vee(\widetilde{e}_jD) + 1 = \varepsilon_j^\vee(D) + 1 = \varepsilon_j^\vee(\widetilde{f}_j^\vee D). \tag{252}$$

2. In the second case we have $\mathrm{jump}_j(\widetilde{e}_jD) \geq 3$, $\mathrm{jump}_j(\widetilde{e}_j\widetilde{f}_j^\vee D) \geq 2$, so that by Proposition 4.17.2c, $\widetilde{e}_j\widetilde{f}_j^\vee\widetilde{f}_j^\vee D \cong \widetilde{f}_j^\vee\widetilde{e}_j\widetilde{f}_j^\vee D \cong \widetilde{f}_j^\vee\widetilde{f}_j^\vee\widetilde{e}_jD$. Then

$$\varepsilon_j^\vee(\widetilde{e}_j\widetilde{f}_j^\vee\widetilde{f}_j^\vee D) = \varepsilon_j^\vee(\widetilde{f}_j^\vee\widetilde{f}_j^\vee\widetilde{e}_jD) = \varepsilon_j^\vee(\widetilde{e}_jD) + 2 = \varepsilon_j^\vee(D) + 2 = \varepsilon_j^\vee(\widetilde{f}_j^\vee\widetilde{f}_j^\vee D). \tag{253}$$

□

We are now ready to prove Theorem 7.10.

Proof. By Theorem 7.1 we have a surjection

$$\mathrm{Ind}\, T_{p;k} \boxtimes \mathcal{R}(A) \twoheadrightarrow A \tag{254}$$

where $k = r(A)$. We will first treat the case where $\mathcal{R}(A) \cong \mathbb{1}$. This implies $A \cong T_{p;k}$, so $\mathbf{0} \neq \widetilde{e}_jA \cong \widetilde{e}_jT_{p;k}$. Note $\varphi_j^{\Lambda_{p(-1)}}(\mathcal{R}(A)) = \delta_{j,p(-1)}$. However, we can never have $\widetilde{e}_{p(-1)}A = \widetilde{e}_{p(-1)}T_{p;k} \neq \mathbf{0}$ as then $\mathrm{Ind}\, T_{p;k-1} \boxtimes L(p(-1)) \twoheadrightarrow A$ has $L(p(-1)) \in \mathrm{Rep}^{\Lambda_{p(-1)}}$ and so $r(A) < k$, violating its minimality. (Recall $|\nu| = k > 1$ in this case.) Therefore $j \neq p(-1)$ and hence $\varphi_j^{\Lambda_{p(-1)}}(\mathcal{R}(A)) = 0$. Thus as $\varepsilon_j(A) = \varepsilon_j(T_{p;k}) > 0 = \varphi_j^{\Lambda_{p(-1)}}(\mathbb{1})$ and we always have trivial surjections

$$\mathrm{Ind}\, T_{p;k} \boxtimes \mathbb{1} \twoheadrightarrow A \quad \text{and} \quad \mathrm{Ind}\, \widetilde{e}_jT_{p;k} \boxtimes \mathbb{1} \twoheadrightarrow \widetilde{e}_jA, \tag{255}$$

(248) always holds and the theorem holds in this case.

For the rest of the proof assume $\mathcal{R}(A) \neq \mathbb{1}$. We divide the proof into the three possible values of $\varepsilon_j(T_{p;k})$: 0, 1, or 2.

- *Case 1*: $\varepsilon_j(T_{p;k}) = 0$

  Since $\varepsilon_j(A) \neq 0$, by Proposition 4.10 there is a surjection

  $$\operatorname{Ind} T_{p;k} \boxtimes \widetilde{e}_j \mathcal{R}(A) \twoheadrightarrow \widetilde{e}_j A. \tag{256}$$

  Note $0 \leq \varphi_j^{\Lambda_{p(-1)}}(\mathcal{R}(A))$ always.

- *Case 2*: $\varepsilon_j(T_{p;k}) = 1$

  - *Case 2a*: $\varphi_j^{\Lambda_{p(-1)}}(\mathcal{R}(A)) = 0 < 1 = \varepsilon_j(T_{p;k})$

    Since $\mathcal{R}(A) \neq \mathbb{1}$, Remark 5.3 implies $\operatorname{jump}_j(\mathcal{R}(A)) = 0$. Because $\varepsilon_j(T_{p;k}) = 1$, then $k \geq 1$ and therefore step $(k-1)$ of the algorithm in Theorem 7.1 provides a surjection

    $$\operatorname{Ind} T_{p;k-1} \boxtimes \widetilde{f}_j^{\vee} \mathcal{R}(A) \twoheadrightarrow A. \tag{257}$$

    If j is part of a class $\mathcal{D}$ pair (j, j') and $\varepsilon(T_{p;k}) = \Lambda_j + \Lambda_{j'}$ then by Proposition 7.7 we can assume that $p(k-1) = j$, $p(k-2) = j'$ so that $\widetilde{e}_j T_{p;k} = T_{p;k-1}$ and $\widetilde{e}_j T_{p;k-1} = \widetilde{e}_j^2 T_{p;k} = \mathbf{0}$. By Proposition 4.10.1b, as $\varepsilon_j(T_{p;k-1}) = 0$, we have

    $$\operatorname{Ind} T_{p;k-1} \boxtimes \widetilde{e}_j \widetilde{f}_j^{\vee} \mathcal{R}(A) \twoheadrightarrow \widetilde{e}_j A. \tag{258}$$

    Because $\operatorname{jump}_j(\mathcal{R}(A)) = 0$, by Proposition 4.14 $\widetilde{f}_j^{\vee} \mathcal{R}(A) \cong \widetilde{f}_j \mathcal{R}(A)$, and so $\widetilde{e}_j \widetilde{f}_j^{\vee} \mathcal{R}(A) \cong \mathcal{R}(A)$. Since $T_{p;k-1} \cong \widetilde{e}_j T_{p;k}$ this gives (248).

  - *Case 2b*: $\varphi_j^{\Lambda_{p(-1)}}(\mathcal{R}(A)) \geq 1 = \varepsilon_j(T_{p;k})$

    Again by Remark 5.3 this implies that $\operatorname{jump}_j(\mathcal{R}(A)) \geq 1$ and $\operatorname{jump}_j(\widetilde{e}_j \mathcal{R}(A)) \geq 2$. Note $\widetilde{e}_j \mathcal{R}(A) \neq \mathbf{0}$ because $\operatorname{jump}_j(\mathcal{R}(A)) \geq 1$ and (257) still holds as well as $\varepsilon_j(T_{p;k-1}) = 0$.

    If $k = 1$, then $T_{p;k} = L(i)$ which forces $j = i$ as $\varepsilon_j(L(i)) = \delta_{ij}$. Further $\mathcal{R}(A) = \widetilde{e}_j^{\vee} A$. We have

    $$\operatorname{Ind} L(j) \boxtimes \widetilde{e}_j^{\vee} A \twoheadrightarrow A. \tag{259}$$

    Because $\operatorname{jump}_j(\widetilde{e}_j^{\vee} A) = \operatorname{jump}_j(\mathcal{R}(A)) \geq 1$ then $\operatorname{jump}_j(\widetilde{e}_j \widetilde{e}_j^{\vee} A) \geq 2$. Proposition 4.17.2c then implies that $\widetilde{e}_j A \cong \widetilde{e}_j \widetilde{f}_i^{\vee} \widetilde{e}_i^{\vee} A \cong \widetilde{e}_j \widetilde{f}_j^{\vee} \widetilde{e}_j^{\vee} A \cong \widetilde{f}_j^{\vee} \widetilde{e}_j \widetilde{e}_j^{\vee} A$. So there is a surjection

    $$\begin{tikzcd} \operatorname{Ind} L(j) \boxtimes \widetilde{e}_j \widetilde{e}_j^{\vee} A \arrow[r, two heads] \arrow[d, phantom, "\wr\|"] & \widetilde{e}_j A \\ \operatorname{Ind} T_{p;1} \boxtimes \widetilde{e}_j \mathcal{R}(A) & \end{tikzcd}$$

    and (249) holds.

    Now assume that $k > 1$ so that by the algorithm in Theorem 7.1, there is a surjection

    $$\operatorname{Ind} T_{p;k-2} \boxtimes \widetilde{f}_{p(k-2)}^{\vee} \widetilde{f}_j^{\vee} \mathcal{R}(A) \twoheadrightarrow A. \tag{260}$$

    Note in Case 2 that $p(k-2) \neq j$ and $\varepsilon_j(T_{p;k-2}) = 0$ and therefore by Proposition 4.10

    $$\operatorname{Ind} T_{p;k-2} \boxtimes \widetilde{e}_j \widetilde{f}_{p(k-2)}^{\vee} \widetilde{f}_j^{\vee} \mathcal{R}(A) \twoheadrightarrow \widetilde{e}_j A. \tag{261}$$

Furthermore since $\mathrm{jump}_j(\widetilde{e}_j\mathcal{R}(A)) \geq 2$, then by Proposition 4.17.1c and 4.17.2c

$$\widetilde{e}_j\widetilde{f}^\vee_{p(k-2)}\widetilde{f}^\vee_j\mathcal{R}(A) \cong \widetilde{f}^\vee_{p(k-2)}\widetilde{e}_j\widetilde{f}^\vee_j\mathcal{R}(A) \cong \widetilde{f}^\vee_{p(k-2)}\widetilde{f}^\vee_j\widetilde{e}_j\mathcal{R}(A). \tag{262}$$

This gives us

$$\operatorname{Ind} T_{p;k-2} \boxtimes \widetilde{f}^\vee_{p(k-2)}\widetilde{f}^\vee_j\widetilde{e}_j\mathcal{R}(A) \twoheadrightarrow \widetilde{e}_jA. \tag{263}$$

By Lemma 7.15,

$$\varepsilon^\vee_{p(-1)}(\widetilde{f}^\vee_j\widetilde{e}_j\mathcal{R}(A)) = \varepsilon^\vee_{p(-1)}(\widetilde{e}_j\widetilde{f}^\vee_j\mathcal{R}(A)) = \varepsilon^\vee_{p(-1)}(\widetilde{f}^\vee_j\mathcal{R}(A)), \tag{264}$$

and $\widetilde{e}_j\mathcal{R}(A) \in \mathrm{Rep}^{\Lambda_i}$. By the minimality of $r(A)$, $\widetilde{f}^\vee_j\mathcal{R}(A), \widetilde{f}^\vee_{p(k-2)}\widetilde{f}^\vee_j\mathcal{R}(A) \notin \mathrm{Rep}^{\Lambda_{p(-1)}}$, hence also considering jump_j, $\widetilde{e}_j(\widetilde{f}^\vee_j\mathcal{R}(A)) = \widetilde{f}^\vee_j\widetilde{e}_j\mathcal{R}(A) \notin \mathrm{Rep}^{\Lambda_{p(-1)}}$ and $\widetilde{f}^\vee_{p(k-2)}\widetilde{f}^\vee_j\widetilde{e}_j\mathcal{R}(A) \notin \mathrm{Rep}^{\Lambda_{p(-1)}}$. When $j, p(k-2) \neq p(-1)$ then $\varepsilon^\vee_{p(k-2)}(\widetilde{f}^\vee_{p(k-2)}\widetilde{f}^\vee_j\widetilde{e}_j\mathcal{R}(A)) = 1 = \varepsilon^\vee_j(\widetilde{f}^\vee_j\widetilde{e}_j\mathcal{R}(A))$ so that applying Lemma 7.4 twice gives

$$\operatorname{Ind} T_{p;k} \boxtimes \widetilde{e}_j\mathcal{R}(A) \twoheadrightarrow \widetilde{e}_jA, \tag{265}$$

agreeing with (249) of Theorem 7.10, which we are in the process of proving.

When $p(k-2) = p(-1)$, then $j \neq p(-1)$. By the minimality of $r(A)$ and Lemma 7.3, it follows in this case that $\varepsilon^\vee_{p(-1)}(\widetilde{f}^\vee_{p(-1)}\widetilde{f}^\vee_j\mathcal{R}(A)) = 2 = \varphi_{p(-1)}(p; k-2)$. By Lemma 7.13 and (262) then $\varepsilon^\vee_{p(-1)}(\widetilde{f}^\vee_{p(-1)}\widetilde{f}^\vee_j\widetilde{e}_j\mathcal{R}(A)) = 2$ so that again by Lemma 7.4 applied to (263) we get

$$\operatorname{Ind} T_{p;k-1} \boxtimes \widetilde{f}^\vee_j\widetilde{e}_j\mathcal{R}(A) \twoheadrightarrow \widetilde{e}_jA \tag{266}$$

and another application of Lemma 7.4 gives (249).

The last case to consider of Case 2b is when $j = p(-1)$. Then $p(k-2) \neq p(-1)$ since $\varepsilon_j(T_{p;k}) = 1$. Using Lemma 7.4 gives (266) as above. Then we either have $\varepsilon^\vee_{p(-1)}(\widetilde{f}^\vee_{p(-1)}\widetilde{e}_{p(-1)}\mathcal{R}(A)) = 2 = \varphi_{p(-1)}(p; k-1)$ if $p(-1)$ is class $\mathcal{A}$ or class $\mathcal{D}$, or $\varepsilon^\vee_{p(-1)}(\widetilde{f}^\vee_{p(-1)}\widetilde{e}_{p(-1)}\mathcal{R}(A)) = 2$ and $\varphi_{p(-1)}(p; k-1) = 3$ when $p(-1)$ is class $\mathcal{B}$. In the former case we can apply Lemma 7.4 to (266), and in the latter case apply Lemma 7.6 to (266) and (263), to get the desired surjection (249).

- *Case 3*: $\varepsilon_j(T_{p;k}) = 2$

  Note that in this case j must be class $\mathcal{B}$ and $p(k-3) \neq j$.

  Note $k > 2$ because p is a cyclotomic path (if $k = 2$ we would have $p(0) = p(1)$). From Theorem 7.1 we have a surjection

$$\operatorname{Ind} T_{p;k-3} \boxtimes \widetilde{f}^\vee_{p(k-3)}\widetilde{f}^\vee_j\widetilde{f}^\vee_j\mathcal{R}(A) \twoheadrightarrow A. \tag{267}$$

  By Proposition 4.10.2 we can apply $\widetilde{e}_j$ to (267) to get

$$\operatorname{Ind} T_{p;k-3} \boxtimes \widetilde{e}_j\widetilde{f}^\vee_{p(k-3)}\widetilde{f}^\vee_j\widetilde{f}^\vee_j\mathcal{R}(A) \twoheadrightarrow \widetilde{e}_jA. \tag{268}$$

- *Case* 3a: $\varphi_j^{\Lambda_{p(-1)}}(\mathcal{R}(A)) < 2 = \varepsilon_j(T_{p;k})$

  Then $\mathrm{jump}_j(\mathcal{R}(A)) \leq 1$ and by formula (117), $\mathrm{jump}_j(\widetilde{f}_j\mathcal{R}(A)) = 0$. Hence by Proposition 4.14.2, $\widetilde{f}_j\widetilde{f}_j\mathcal{R}(A) \cong \widetilde{f}_j^\vee\widetilde{f}_j\mathcal{R}(A)$. Furthermore because $p(k-3) \neq j$, by Proposition 4.17.1c, $\widetilde{e}_j$ and $\widetilde{f}^\vee_{p(k-3)}$ commute. Thus we have

  $$\widetilde{e}_j\widetilde{f}^\vee_{p(k-3)}\widetilde{f}^\vee_j\widetilde{f}^\vee_j\mathcal{R}(A) \cong \widetilde{f}^\vee_{p(k-3)}\widetilde{e}_j\widetilde{f}^\vee_j\widetilde{f}^\vee_j\mathcal{R}(A) \cong \widetilde{f}^\vee_{p(k-3)}\widetilde{e}_j\widetilde{f}_j\widetilde{f}^\vee_j\mathcal{R}(A) \cong \widetilde{f}^\vee_{p(k-3)}\widetilde{f}^\vee_j\mathcal{R}(A). \tag{269}$$

  The map in (268) then becomes

  $$\operatorname{Ind} T_{p;k-3} \boxtimes \widetilde{f}^\vee_{p(k-3)}\widetilde{f}^\vee_j\mathcal{R}(A) \twoheadrightarrow \widetilde{e}_j A. \tag{270}$$

  Note that since j is class $\mathcal{B}$ then $p(k-3)$ must be class $\mathcal{A}$. If $p(k-3), j \neq p(-1)$, then $\varepsilon^\vee_{p(k-3)}(\widetilde{f}^\vee_{p(k-3)}\widetilde{f}^\vee_j\mathcal{R}(A)) = 1$ and $\varepsilon^\vee_j(\widetilde{f}^\vee_j\mathcal{R}(A)) = 1$ so that two applications of Lemma 7.4 give

  $$\begin{tikzcd} \operatorname{Ind} T_{p;k-1} \boxtimes \mathcal{R}(A) \arrow[r, two heads] \arrow[d, phantom, "\wr\|"] & \widetilde{e}_j A \\ \widetilde{e}_j T_{p;k} & \end{tikzcd} \tag{271}$$

  yielding (248) of Theorem 7.10.

  If $p(k-3) = p(-1)$ then by Lemma 7.3 and the Serre relations, $\varepsilon^\vee_{p(k-3)}(\widetilde{f}^\vee_{p(k-3)}\widetilde{f}^\vee_j\mathcal{R}(A)) = 2 = \varphi_{p(k-3)}(p;k-3)$. Then we may again apply Lemma 7.4 twice to (270) to get (271).

  Finally, in the case where $j = p(-1)$ then because $p(k-3) \neq j = p(-1)$ we can again apply Lemma 7.4 to (270) to get

  $$\operatorname{Ind} T_{p;k-2} \boxtimes \widetilde{f}^\vee_j\mathcal{R}(A) \twoheadrightarrow \widetilde{e}_j A. \tag{272}$$

  Then $\varepsilon^\vee_{p(-1)}(\widetilde{f}^\vee_j\mathcal{R}(A)) = \varepsilon^\vee_{p(-1)}(\widetilde{f}^\vee_{p(-1)}\mathcal{R}(A)) = 2$ and $\varphi_j(p;k-2) = 3$ so that by Lemma 7.6 a surjection like (271) exists. (Recall the definition of cyclotomic path does not allow $\varphi_j(p;k-2) = 4$.)

- *Case* 3b: $\varphi_j^{\Lambda_{p(-1)}}(\mathcal{R}(A)) \geq 2 = \varepsilon_j(T_{p;k})$

  Then $\mathrm{jump}_j(\mathcal{R}(A)) \geq 2$. This implies $\widetilde{e}_j\mathcal{R}(A) \neq \mathbf{0}$ by (267). It also follows that $\mathrm{jump}_j(\widetilde{f}^\vee_j\mathcal{R}(A)) \geq 1$, $\mathrm{jump}_j(\widetilde{e}_j\widetilde{f}^\vee_j\mathcal{R}(A)) \geq 2$ so that by Proposition 4.17.2c, $\widetilde{e}_j\widetilde{f}^\vee_j\widetilde{f}^\vee_j\mathcal{R}(A) \cong \widetilde{f}^\vee_j\widetilde{e}_j\widetilde{f}^\vee_j\mathcal{R}(A)$. Also, $\mathrm{jump}_j(\widetilde{e}_j\mathcal{R}(A)) \geq 3$ so again by Proposition 4.17.2c $\widetilde{e}_j\widetilde{f}^\vee_j\mathcal{R}(A) \cong \widetilde{f}^\vee_j\widetilde{e}_j\mathcal{R}(A)$. Again because $p(k-3) \neq j$, by Proposition 4.17.1c, $\widetilde{f}^\vee_{p(k-3)}$ and $\widetilde{e}_j$ commute. Together these imply

  $$\widetilde{e}_j\widetilde{f}^\vee_{p(k-3)}\widetilde{f}^\vee_j\widetilde{f}^\vee_j\mathcal{R}(A) \cong \widetilde{f}^\vee_{p(k-3)}\widetilde{e}_j\widetilde{f}^\vee_j\widetilde{f}^\vee_j\mathcal{R}(A) \cong \widetilde{f}^\vee_{p(k-3)}\widetilde{f}^\vee_j\widetilde{f}^\vee_j\widetilde{e}_j\mathcal{R}(A). \tag{273}$$

  Considering cases similar to those in Case 3a, applying Lemma 7.4 and Lemma 7.6 to (268), we get

  $$\operatorname{Ind} T_{p;k} \boxtimes \widetilde{e}_j\mathcal{R}(A) \twoheadrightarrow \widetilde{e}_j A \tag{274}$$

  agreeing with (249).

This concludes the proof of Theorem 7.10 apart from observing that in the case $p(k-3) = p(-1) \neq j$ it is useful to note

$$\varepsilon^{\vee}_{p(-1)}(\widetilde{f}^{\vee}_{p(k-1)}\widetilde{f}^{\vee}_{j}\widetilde{f}^{\vee}_{j}\widetilde{e}_j\mathcal{R}(A)) = \varepsilon^{\vee}_{p(-1)}(\widetilde{f}^{\vee}_{p(k-1)}\widetilde{f}^{\vee}_{j}\widetilde{f}^{\vee}_{j}\mathcal{R}(A)) \tag{275}$$

by Lemma 7.3 and Lemma 7.15.

□

Example 7.16. Recall that in type $C^{(1)}_{\ell}$, $B^{1,1} \otimes B(\Lambda_1) \simeq B(\Lambda_0) \oplus B(\Lambda_2)$. We illustrate the module-theoretic phenomena that mirrors this fact. It holds that

$$L(1) \in \mathrm{Rep}^{\Lambda_1} \qquad \widetilde{f}^{\vee}_0 L(1) \simeq \widetilde{f}_1 L(0) \simeq \mathcal{L}(01) \in \mathrm{Rep}^{\Lambda_0} \qquad \widetilde{f}^{\vee}_2 L(1) \simeq \mathcal{L}(21) \in \mathrm{Rep}^{\Lambda_2}\,. \tag{276}$$

In other words, modules A both from Rep^{Λ_0} and Rep^{Λ_2} can lead to modules $\mathcal{R}(A) \in \mathrm{Rep}^{\Lambda_1}$ when one performs the algorithm of Theorem 7.1. Regarding Example 2.2, Remark 7.9, and Theorem 7.10, this corresponds to the fact that

$$\widetilde{f}_1\widetilde{f}_0 b_{\Lambda_0} = \mathcal{T}(\widetilde{f}_1\widetilde{f}_0(\boxed{\bar{1}} \otimes b_{\Lambda_1})) = \mathcal{T}(\boxed{1} \otimes b_{\Lambda_1}) \tag{277}$$

whereas

$$\widetilde{f}_1\widetilde{f}_2 b_{\Lambda_2} = \mathcal{T}(\widetilde{f}_1\widetilde{f}_2(\boxed{1} \otimes b_{\Lambda_1})) = \mathcal{T}(\boxed{2} \otimes \widetilde{f}_1 b_{\Lambda_1}). \tag{278}$$

A more trivial example of the same phenomenon is given by $\mathbb{1} \in \mathrm{Rep}^{\Lambda_1}$, $\widetilde{f}^{\vee}_0\,\mathbb{1} \simeq L(0) \in \mathrm{Rep}^{\Lambda_0}$, $\widetilde{f}^{\vee}_2\,\mathbb{1} \simeq L(2) \in \mathrm{Rep}^{\Lambda_2}$. This corresponds to $\widetilde{f}_0 b_{\Lambda_0} = \mathcal{T}(\widetilde{f}_0(\boxed{\bar{1}} \otimes b_{\Lambda_1})) = \mathcal{T}(\boxed{0} \otimes b_{\Lambda_1})$ and $\widetilde{f}_2 b_{\Lambda_2} = \mathcal{T}(\widetilde{f}_2(\boxed{1} \otimes b_{\Lambda_1})) = \mathcal{T}(\boxed{2} \otimes b_{\Lambda_1})$.

8 Appendix

8.1 Simple modules in $\mathrm{Rep}^{\Lambda_h+\Lambda_i}$ in rank 2

In Figure 3 we draw the highest weight crystal $B(\Lambda_h + \Lambda_i)$ in type B_2/C_2. That is, the associated Cartan matrix is

$$\begin{pmatrix} 2 & -2 \\ -1 & 2 \end{pmatrix} \tag{279}$$

with $a_{hi} = -1$ and $a_{ih} = -2$.

In Figure 4 we redraw this crystal with nodes the corresponding simple $R(m_h\alpha_h + m_i\alpha_i)$-modules in $\mathrm{Rep}^{\Lambda_h+\Lambda_i}$, $m_h, m_i \geq 0$. The cyclotomic condition $\mathcal{I}^{\Lambda_h+\Lambda_i}M = \mathbf{0}$ forces $m_h + m_i \leq 7$. Indeed we can see this from Figure 3 or Figure 4.

In Figure 4 we indicate $\mathrm{jump}_i(M), \mathrm{jump}_h(M)$, in part so one can see when $\widetilde{f}_* M \in \mathrm{Rep}^{\Lambda_i+\Lambda_h}$. Furthermore when $\mathrm{jump}_h(M) = 0$ we draw $M \xrightarrow{\widetilde{f}_h} \mathrm{Ind}\, M \boxtimes L(h)$ (similarly for i) to mark the fact $\widetilde{f}_h M \cong \widetilde{f}^{\vee}_h M$ is irreducibly induced. In other cases when modules can be realized as induced from other modules (without needing to take cosocle) we also express the module in this form. The only exception to this is the 2-dimensional module $\widetilde{f}_h\mathcal{L}(hii)$, whose character has support $[hiih]$.



Figure 3: Crystal graph $B(\Lambda_h + \Lambda_i)$ in type B_2.

References

[1] Susumu Ariki and Kazuhiko Koike, *A Hecke algebra of* $(\mathbf{Z}/r\mathbf{Z}) \wr \mathfrak{S}_n$ *and construction of its irreducible representations*, Adv. Math. **106** (1994), no. 2, 216–243.

[2] Georgia Benkart, Seok-Jin Kang, Se-jin Oh, and Euiyong Park, *Construction of irreducible representations over Khovanov-Lauda-Rouquier algebras of finite classical type*, Int. Math. Res. Not. IMRN (2014), no. 5, 1312–1366. MR 3178600

[3] Michel Broué and Gunter Malle, *Zyklotomische Heckealgebren*, Astérisque (1993), no. 212, 119–189, Représentations unipotentes génériques et blocs des groupes réductifs finis.

[4] Jonathan Brundan and Alexander Kleshchev, *Blocks of cyclotomic Hecke algebras and Khovanov-Lauda algebras*, Invent. Math. **178** (2009), no. 3, 451–484.

[5] ______, *Graded decomposition numbers for cyclotomic Hecke algebras*, Adv. Math. **222** (2009), no. 6, 1883–1942.

[6] I. V. Cherednik, *A new interpretation of Gel′fand-Tzetlin bases*, Duke Math. J. **54** (1987), no. 2, 563–577.

[7] Ghislain Fourier, Masato Okado, and Anne Schilling, *Perfectness of Kirillov-Reshetikhin crystals for nonexceptional types*, Quantum affine algebras, extended affine Lie algebras, and their applications, Contemp. Math., vol. 506, Amer. Math. Soc., Providence, RI, 2010, pp. 127–143.

[8] I. Grojnowski, *Affine sl_p controls the representation theory of the symmetric group and related Hecke algebras*, arXiv:9907129 (1999).



Figure 4: The crystal graph $B(\Lambda_h + \Lambda_i)$ in type B_2 with nodes labeled by corresponding simple modules in $\text{Rep}^{\Lambda_h+\Lambda_i}$ and jump indicated. When $\text{jump}_i(M) = 0$ ($\text{jump}_h(M) = 0$) we indicate with a thick i-arrow (respectively h-arrow).

[9] Victor G. Kac, *Infinite-dimensional Lie algebras*, second ed., Cambridge University Press, Cambridge, 1985.

[10] Seok-Jin Kang and Masaki Kashiwara, *Quantized affine algebras and crystals with core*, Comm. Math. Phys. **195** (1998), no. 3, 725–740.

[11] ———, *Categorification of highest weight modules via Khovanov-Lauda-Rouquier algebras*, Invent. Math. **190** (2012), no. 3, 699–742.

[12] Seok-Jin Kang, Masaki Kashiwara, Kailash C. Misra, Tetsuji Miwa, Toshiki Nakashima, and Atsushi Nakayashiki, *Affine crystals and vertex models*, Infinite analysis, Part A, B (Kyoto, 1991), Adv. Ser. Math. Phys., vol. 16, World Sci. Publ., River Edge, NJ, 1992, pp. 449–484.

[13] ———, *Perfect crystals of quantum affine Lie algebras*, Duke Math. J. **68** (1992), no. 3, 499–607.

[14] M. Kashiwara, *On crystal bases of the Q-analogue of universal enveloping algebras*, Duke Math. J. **63** (1991), no. 2, 465–516.

[15] Masaki Kashiwara, *Bases cristallines*, C. R. Acad. Sci. Paris Sér. I Math. **311** (1990), no. 6, 277–280.

[16] ———, *On crystal bases*, Representations of groups (Banff, AB, 1994), CMS Conf. Proc., vol. 16, Amer. Math. Soc., Providence, RI, 1995, pp. 155–197.

[17] Masaki Kashiwara and Yoshihisa Saito, *Geometric construction of crystal bases*, Duke Math. J. **89** (1997), no. 1, 9–36.

[18] Mikhail Khovanov and Aaron D. Lauda, *A diagrammatic approach to categorification of quantum groups. I*, Represent. Theory **13** (2009), 309–347.

[19] ———, *A diagrammatic approach to categorification of quantum groups II*, Trans. Amer. Math. Soc. **363** (2011), no. 5, 2685–2700.

[20] A. N. Kirillov and N. Yu. Reshetikhin, *Representations of Yangians and multiplicities of the inclusion of the irreducible components of the tensor product of representations of simple Lie algebras*, Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI) **160** (1987), no. Anal. Teor. Chisel i Teor. Funktsii. 8, 211–221, 301.

[21] A. Kleshchev, *Modular Representation Theory of Symmetric Groups*, arXiv:1405.3326 (2014).

[22] A. S. Kleshchev, *Branching rules for modular representations of symmetric groups. I*, J. Algebra **178** (1995), no. 2, 493–511.

[23] Aaron D. Lauda and Monica Vazirani, *Crystals from categorified quantum groups*, Adv. Math. **228** (2011), no. 2, 803–861.

[24] Cristian Lenart and Anne Schilling, *Crystal energy functions via the charge in types A and C*, Math. Z. **273** (2013), no. 1-2, 401–426.

[25] Ivan Losev and Ben Webster, *On uniqueness of tensor products of irreducible categorifications*, Selecta Math. (N.S.) **21** (2015), no. 2, 345–377. MR 3338680

[26] George Lusztig, *Introduction to quantum groups*, Modern Birkhäuser Classics, Birkhäuser / Springer, New York, 2010, Reprint of the 1994 edition.

[27] Masato Okado and Anne Schilling, *Existence of Kirillov-Reshetikhin crystals for nonexceptional types*, Represent. Theory **12** (2008), 186–207.

[28] M. Vazirani, *A Hecke theoretic shadow of tensoring the crystal of the basic representation with a level 1 perfect crystal*, (1999), Unpublished notes.

[29] ———, *Irreducible modules over the affine Hecke algebra: a strong multiplicity one result*, Ph.D. thesis, UC Berkeley, 1999.

[30] ———, *Filtrations on the Mackey decomposition for cyclotomic Hecke algebras*, J. Algebra **252** (2002), no. 2, 205–227.

[31] ———, *A Hecke theoretic shadow of tensoring the crystal of the basic representation with a level 1 perfect crystal*, Mathematisches Forschungsinstitut Oberwolfach Report **Report No. 14/2003** (2003).

[32] ———, *An observation on highest weight crystals*, J. Algebra **315** (2007), no. 2, 483–501.

[33] ———, *Categorifying the tensor product of a level 1 highest weight and perfect crystal in type A*, Proceedings of the 2012-2014 Southeastern Lie Theory Workshops, Proc. Sympos. Pure Math., Amer. Math. Soc., Providence, RI, To appear, arXiv.

[34] B. Webster, *Knot invariants and higher representation theory I: diagrammatic and geometric categorification of tensor products*, arXiv:1001.2020 (2010).

[35] ———, *Knot invariants and higher representation theory*, Memoirs of the AMS (to appear).